

US-FT/13-94
 (Revised version)
 hep-th/9409016
 August 1994

$\mathcal{W}$ -ALGEBRAS FROM SOLITON EQUATIONS AND HEISENBERG SUBALGEBRAS

CARLOS R. FERNÁNDEZ-POUSA¹, MANUEL V. GALLAS²,
 J. LUIS MIRAMONTES³, and JOAQUÍN SÁNCHEZ GUILLÉN⁴

*Departamento de Física de Partículas,
 Facultad de Física,
 Universidad de Santiago,
 E-15706 Santiago de Compostela, Spain*

ABSTRACT

We derive sufficient conditions under which the “second” Hamiltonian structure of a class of generalized KdV-hierarchies defines one of the classical $\mathcal{W}$ -algebras obtained through Drinfel’d-Sokolov Hamiltonian reduction. These integrable hierarchies are associated to the Heisenberg subalgebras of an untwisted affine Kac-Moody algebra. When the principal Heisenberg subalgebra is chosen, the well known connection between the Hamiltonian structure of the generalized Drinfel’d-Sokolov hierarchies —the Gel’fand-Dickey algebras— and the $\mathcal{W}$ -algebras associated to the Casimir invariants of a Lie algebra is recovered. After carefully discussing the relations between the embeddings of $A_1 = sl(2, \mathbb{C})$ into a simple Lie algebra g and the elements of the Heisenberg subalgebras of $g^{(1)}$, we identify the class of $\mathcal{W}$ -algebras that can be defined in this way. For A_n , this class only includes those

¹ pouasa@gaes.usc.es

² gallas@gaes.usc.es

³ miramontes@gaes.usc.es

⁴ joaquin@gaes.usc.es

associated to the embeddings labelled by partitions of the form $n + 1 = k(m) + q(1)$ and $n + 1 = k(m + 1) + k(m) + q(1)$.

1. Introduction.

The recent interest in $\mathcal{W}$ -algebras is motivated by their wide spectrum of applications in classical and quantum two-dimensional field theory. Just to mention some of them, $\mathcal{W}$ -algebras play an important role in the study of conformal field theories, quantum and topological gravity, string theories, critical phenomena in statistical mechanics, and even the quantum Hall effect (see the reviews [1]). Moreover, from a fundamental point of view, they might lead to the formulation of Lorentz-invariant field theories with symmetries generated by fields of higher spin.

The $\mathcal{W}$ -algebras are (non-linear) extensions of the Virasoro algebra by conformal primary fields [2]. The algebraic structures related to $\mathcal{W}$ -algebras first appeared in the context of certain integrable hierarchies of differential equations [3]. The simplest example of this relation is provided by the Korteweg-de Vries equation (KdV), the archetypical integrable system,

$$\begin{aligned}\partial_t u(x, t) &= -\frac{1}{4}u'''(x, t) + \frac{3}{2}u(x, t)u'(x, t) \\ &= \{H, u(x, t)\}_2,\end{aligned}$$

where $' = \partial/\partial x$, the Hamiltonian is $H = \int dx \frac{1}{4}u^2(x, t)$, and the “second” Poisson bracket is

$$\{u(x, t), u(y, t)\}_2 = \frac{1}{2}\delta'''(x - y) - 2u(x, t)\delta'(x - y) - u'(x, t)\delta(x - y),$$

which is nothing but a classical realization of the Virasoro algebra. In [4], it was shown that generalized equations of the KdV type can be defined for arbitrary affine Kac-Moody algebras; the original KdV equation corresponding to $A_1^{(1)}$. The “second” Hamiltonian structure of these integrable differential equations is defined by the Gel’fand-Dickey algebras, and it has been proved [5] that they are connected with the $\mathcal{W}$ -algebras associated to the Casimir invariants of the classical Lie algebras [6]. For instance, the second Hamiltonian structure of the Drinfel’d-Sokolov A_{n-1} generalized KdV hierarchy is connected with $\mathcal{W}_n$, the $\mathcal{W}$ -algebra whose generators have conformal spin $2, 3, \dots, n$ [7].

This early known connection between the Drinfel’d-Sokolov generalized KdV hierarchies and (certain) $\mathcal{W}$ -algebras relates them to Lie algebras, and this opens the possibility of their classification. In fact, among the different methods used in the literature to construct $\mathcal{W}$ -algebras, the most effective and systematic one is conceptually inspired

by this connection. It consists in the Hamiltonian reduction of affine Kac-Moody algebras [1,8,9,10], and it is usually called (quantum or classical) Drinfel'd-Sokolov reduction. With this method, a polynomial extension of the Virasoro algebra can be associated to each embedding of $A_1 \equiv sl(2, \mathbb{C})$ into a simple finite Lie algebra [8,11]. The Lagrangian field theoretical version of this procedure involves the reduction of Wess-Zumino-Novikov-Witten models (WZNW) to, generally non-abelian, Toda field theories, which are well known two-dimensional integrable field theories. Moreover, there is convincing evidence to believe that all the $\mathcal{W}$ -algebras can be constructed in this way [12].

Using the Drinfel'd-Sokolov reduction procedure at the classical level, the $\mathcal{W}$ -algebra associated to the Casimir invariants of the Lie algebra g is recovered by choosing the principal embedding of A_1 into g . The natural question is if it is possible to associate integrable systems to all the other $\mathcal{W}$ -algebras obtained with other embeddings. So far, the outcome of all the attempts to answer this question is the construction of integrable systems associated to some particular families of $\mathcal{W}$ -algebras [13,14,15,16, 17,18,19,20], but still the general structure of this connection has not been developed.

Recently, the Drinfel'd-Sokolov construction has been generalized showing that it is possible to associate an integrable hierarchy of differential equations of the KdV type to each particular Heisenberg subalgebra of an affine Kac-Moody algebra [21,22] (see also [19,23,24,25]). The original Drinfel'd-Sokolov hierarchies are then recovered with the principal Heisenberg subalgebra. Even more, when the affine Kac-Moody algebra is simply-laced, it has been proved in ref. [26] that the hierarchy is one of those constructed by Kac and Wakimoto within the tau-function approach [27]. These integrable hierarchies are bi-Hamiltonian, and one of their Hamiltonian structures, the “second” one, exhibits a conformal invariance [22]. Therefore, it is logical to think that their Hamiltonian structures might be related to the $\mathcal{W}$ -algebras obtained by Hamiltonian reduction of affine Kac-Moody algebras. It is the purpose of this paper to establish the precise form of this connection.

A priori, the approach of refs.[21,22] seems quite different from that of [1,8,9,10]. In the later, the fundamental objects are the embeddings of A_1 into simple Lie algebras, while, in the former, the construction is based on the Heisenberg subalgebras of affine Kac-Moody algebras. Besides, in the Drinfel'd-Sokolov reduction, the $\mathcal{W}$ -algebras are obtained by reducing a classical Kac-Moody algebra that is formulated as the Kirillov-Poisson bracket associated to a Lie algebra g , *i.e.*, g is affinized once in a variable x . In contrast, the

integrable hierarchies of [21,22] are defined in terms of Lax operators defined on a loop algebra $\mathcal{L}(g) = g \otimes \mathbb{C}[z, z^{-1}]$ —an affine Kac-Moody algebra in general; therefore, the phase space consists of functions of a variable x on $\mathcal{L}(g)$, *i.e.*, g is affinized twice: in z and x . However, notice that the Lie algebra g can be identified with the subalgebra $g \otimes 1 \subset \mathcal{L}(g)$, which indicates that the phase space involved in the Drinfel'd-Sokolov reductions is, in fact, a subset of the phase space where the generalized integrable hierarchies are defined. We will show how this relation between these two phase spaces specifies a connection between the embeddings of A_1 into g and the Heisenberg subalgebras of $\mathcal{L}(g)$.

The paper is organized as follows. In Section 2 we review the construction of [21,22], where integrable hierarchies are associated to the non-equivalent Heisenberg subalgebras of a loop algebra. The definition of these integrable systems involves the choice of a constant graded element of a Heisenberg subalgebra, say Λ . A distinction is made between “regular” and “non-regular” Λ 's, leading to type I and type II hierarchies, respectively. Nevertheless, even though it has been proved in [19,25] that most Heisenberg subalgebras do not have regular graded elements, type II hierarchies are not described in detail in the original references. We have tried to fill this gap in Section 2.

In the next two sections we analyse the “second” Hamiltonian structure of the integrable hierarchies of [21,22] and we compare it with the $\mathcal{W}$ -algebras. In Section 3 we discuss the conformal invariance of the second Poisson bracket and we find the energy-momentum tensor that generates this conformal transformation on a subspace of the total phase space. Restricted to this subspace, the Poisson bracket is just the Kirillov-Poisson bracket associated to a (non-simple, in general) Lie algebra, say $\widehat{g}_0(\mathbf{s})$. This is the starting point in connecting the second Poisson bracket algebra with $\mathcal{W}$ -algebras. The phase space of the integrable hierarchy consists of a set of equivalence classes under a group of gauge transformations; we propose a convenient choice of the gauge slice when Λ satisfies a “non-degeneracy” condition. In Section 4 we establish the connection between the second Poisson bracket algebra and the classical $\mathcal{W}$ -algebras obtained with the Drinfel'd-Sokolov reduction approach. This connection, explained in theorem 2, gives a precise relation between the constant graded element Λ used in the construction of the integrable hierarchy and the J_+ of the embedding of A_1 into $\widehat{g}_0(\mathbf{s})$ that specifies the corresponding $\mathcal{W}$ -algebra.

With this result, it is possible to investigate which $\mathcal{W}$ -algebras can be constructed as the second Poisson bracket algebras of the integrable hierarchies of [21,22]. This is equivalent to show which nilpotent elements J_+ are related to the graded elements of

the Heisenberg subalgebras, and we do it here for the classical Lie algebras of the A_n series. Recall that the inequivalent Heisenberg subalgebras of the untwisted Kac-Moody algebra $g^{(1)}$ are in one-to-one relation with the conjugacy classes of the Weyl group of g . Therefore, as a previous step, we have summarized in Section 5 the classification of the A_1 subalgebras and the conjugacy classes of the Weyl group of A_n . In order to establish the required connections between the elements of all the Heisenberg subalgebras of $A_n^{(1)}$ and the A_1 subalgebras of A_n . In fact, we derive two kinds of connections. The first one is based on the properties of the cyclic element of B. Kostant [28] and it can be generalized to the other classical algebras. The second one is based on the particular structure of the Heisenberg subalgebras of $A_n^{(1)}$ and, hence, it is specific of this case. These results are interesting by themselves, and we hope that they will be useful, for instance, in the study of two-dimensional integrable field theories [29,30]. The class of $\mathcal{W}$ -algebras that can be constructed as the second Hamiltonian structures of the integrable hierarchies of [21,22] is obtained in Section 6. This class is quite limited, even if type II hierarchies are included, and the origin of this restriction is that J_+ has to be related to a “graded” element of a Heisenberg subalgebra.

In Section 7 we consider some examples for the purposes of illustration of our results. Finally, we present our conclusions in Section 8. The explicit construction of the inequivalent Heisenberg subalgebras of $A_n^{(1)}$, with $n = 2, \dots, 5$, is presented in the Appendix.

2. The generalized KdV hierarchies.

In this section we summarize the construction of [21,22], and we generalize some of the results of these works to include the case of the hierarchies of type II. Unless otherwise indicated, we follow the conventions of [22].

The central object is an affine Kac-Moody algebra $\hat{g}$. In this paper, $\hat{g}$ will be considered as the loop algebra associated to a finite (simple) Lie algebra g , $\hat{g} = g \otimes \mathbb{C}[z, z^{-1}] \oplus \mathbb{C} d$, where $d = z d/dz$, and whose central extension will be ignored (but see ref.[26]). A very important role is played by the Heisenberg subalgebras of $\hat{g}$, which are maximal abelian subalgebras of $\hat{g}$ (see [31] for a precise definition). Up to conjugation, they are in one-to-one correspondence with the conjugacy classes of the Weyl group of g [31], and we denote them as $\mathcal{H}[w]$, where $[w]$ indicates the conjugacy class. Associated

to $\mathcal{H}[w]$ there is a distinguished gradation $\mathbf{s}_w$, with the property that $\mathcal{H}[w]$ is an invariant subspace under the adjoint action of $d_{\mathbf{s}_w}$ ¹. This ensures that one can choose a basis of $\mathcal{H}[w]$ whose elements are graded with respect to $\mathbf{s}_w$.

The construction of the integrable hierarchies is based on the (matrix) Lax equation. The Lax operator is associated to the data $\{\widehat{g}, \mathcal{H}[w], \mathbf{s}_w, \mathbf{s}, \Lambda\}$, where $\mathbf{s}$ is an additional gradation such that $\mathbf{s} \preceq \mathbf{s}_w$ with respect to a partial ordering, and Λ is a constant element of $\mathcal{H}[w]$ with positive $\mathbf{s}_w$ -grade i . Then, the Lax operator is defined as

$$L = \partial_x + \Lambda + q(x), \quad (2.1)$$

where the potential $q(x)$ is a function of x on the subspace

$$Q = \widehat{g}_{\geq 0}(\mathbf{s}) \cap \widehat{g}_{< i}(\mathbf{s}_w). \quad (2.2)$$

For technical reasons, the potentials are taken to be periodic functions of x [4]. Notice that any element $\Lambda \in \mathcal{H}[w]$ is semisimple, *i.e.*, $\text{ad } \Lambda$ is diagonalizable on $\widehat{g}$; then $\widehat{g} = \text{Ker}(\text{ad } \Lambda) \oplus \text{Im}(\text{ad } \Lambda)$, where $\text{Ker}(\text{ad } \Lambda)$ is a subalgebra of $\widehat{g}$, and $[\text{Ker}(\text{ad } \Lambda), \text{Im}(\text{ad } \Lambda)] \subseteq \text{Im}(\text{ad } \Lambda)$. A distinction is made between hierarchies of type I and type II, referring to whether the element Λ is “regular” or not —regularity implying $\text{Ker}(\text{ad } \Lambda) = \mathcal{H}[w]$.

The potential $q(x)$ plays the role of the phase space coordinate in this system. However, there exist symmetries corresponding to the gauge transformations

$$L \mapsto \exp(\text{ad } S)(L), \quad (2.3)$$

generated by the functions S of x on the subspace

$$P = \widehat{g}_0(\mathbf{s}) \cap \widehat{g}_{< 0}(\mathbf{s}_w). \quad (2.4)$$

The flows of the integrable hierarchy are defined on the set of gauge invariant functionals of $q(x)$. It is straightforward to find a basis for these gauge invariant functionals. One simply performs a non-singular gauge transformation to take $q(x)$ to some canonical form $q^{(\text{can})}(x)$; then, the components of $q^{(\text{can})}(x)$ and their x -derivatives provide the desired basis.

¹ With the exception of a few cases including the principal Heisenberg subalgebras, the choice of $\mathbf{s}_w$ is not unique, something that has not been mentioned in [21,22]. We shall comment on this point later.

The flows that define the hierarchies are non-linear partial differential equations, and it is convenient to demand that they can be written in the form

$$\frac{\partial q^{(\text{can})}}{\partial t} = F \left(q^{(\text{can})}, \partial_x q^{(\text{can})}, \dots \right), \quad (2.5)$$

with F being a polynomial functional of the components of $q^{(\text{can})}$ and its x -derivatives; otherwise one would obtain some sort of “rational” non-linear differential equations. The condition that the map $\text{ad } \Lambda : P \mapsto Q$ is injective ensures this polynomiality; it is equivalent to

$$\text{Ker}(\text{ad } \Lambda) \cap P = \emptyset, \quad (2.6)$$

and it allows the use of the standard Drinfel’d-Sokolov gauge fixing procedure. Let us recall that, within this procedure, $q^{(\text{can})}(x)$ is characterized as an element of some complementary vector space of $[\Lambda, P]$ in Q , *i.e.*, $q^{(\text{can})}(x) \in Q^{(\text{can})}$ with $Q = Q^{(\text{can})} + [\Lambda, P]$. The choice of $Q^{(\text{can})}$ is not unique (for a very detailed description of the Drinfel’d-Sokolov gauge fixing in a similar situation, see [12]).

If the hierarchy is of type I, then $\text{Ker}(\text{ad } \Lambda) \cap P = \emptyset$ for any choice of Λ because of the following.

Lemma 1. *Let $b \in \mathcal{H}[w]$ be an element of $\mathcal{H}[w]$ that is graded with respect to the two gradations $\mathbf{s}$ and $\mathbf{s}'$. Then, the $\mathbf{s}$ -grade of b equals $N_{\mathbf{s}}/N_{\mathbf{s}'}$ times its $\mathbf{s}'$ -grade.*

Proof. Recall that the derivation inducing the gradation $\mathbf{s}$ is

$$d_{\mathbf{s}} = N_{\mathbf{s}} z \frac{d}{dz} + H_{\mathbf{s}}. \quad (2.7)$$

$H_{\mathbf{s}}$ is an element of the Cartan subalgebra of g such that $\alpha_j(H_{\mathbf{s}}) = s_j \in \mathbb{Z}^+$, where $\alpha_1, \dots, \alpha_{\text{rank}(g)}$ is a choice of the simple roots of g , and

$$N_{\mathbf{s}} = \sum_{j=0}^{\text{rank}(g)} k_j s_j, \quad (2.8)$$

with $\{k_0, \dots, k_{\text{rank}(g)}\}$ being the Kac labels of $\widehat{g}$ (to compare with [22], notice that $H_{\mathbf{s}}/N_{\mathbf{s}} \equiv \delta_{\mathbf{s}} \cdot H$). Therefore,

$$\frac{d_{\mathbf{s}}}{N_{\mathbf{s}}} - \frac{d_{\mathbf{s}'}}{N_{\mathbf{s}'}} = \frac{H_{\mathbf{s}}}{N_{\mathbf{s}}} - \frac{H_{\mathbf{s}'}}{N_{\mathbf{s}'}} \equiv \Delta(\mathbf{s}, \mathbf{s}') \quad (2.9)$$

is also an element of the Cartan subalgebra of g . If $b \in \mathcal{H}[w]$ is graded with respect to the two gradations, then, it has to be stable under the adjoint action of $\Delta(\mathbf{s}, \mathbf{s}')$, which means that $[\Delta(\mathbf{s}, \mathbf{s}'), b] = \beta b$ for some rational number β . Now, taking inner products with the elements of $\mathcal{H}[w]$ and using the invariance of the Killing form on $\widehat{g}$ and the commutativity of $\mathcal{H}[w]$, it follows that $\beta = 0$. Consequently,

$$[d_{\mathbf{s}}, b] = \frac{N_{\mathbf{s}}}{N_{\mathbf{s}'}} [d_{\mathbf{s}'}, b] \quad (2.10)$$

which proves the lemma. $\square$

On the contrary, if the hierarchy is of type II one has to restrict the choice of Λ in such a way that $\text{Ker}(\text{ad } \Lambda) \cap P = \emptyset$ to ensure the polynomiality of the flows.

The outcome of [21,22] is that there exists an infinite number of commuting flows on the gauge equivalence classes of L . These flows have the following form. For each constant element b of the centre of $\text{Ker}(\text{ad } \Lambda)$ with non-negative $\mathbf{s}_w$ -grade there is a flow that can be defined in many gauge equivalent ways parameterized by the generators $\theta(x) \in P$ of (infinitesimal) gauge transformations²:

$$\begin{aligned} \frac{\partial q(x)}{\partial t_b^\theta} &= [\text{P}_{\geq 0[\mathbf{s}_w]} (\exp(-\text{ad } V) (b)) + \theta, L] \\ &= [-\text{P}_{< 0[\mathbf{s}_w]} (\exp(-\text{ad } V) (b)) + \theta, L]; \end{aligned} \quad (2.11)$$

notice that the centre of $\text{Ker}(\text{ad } \Lambda)$, $\text{Cent}(\text{Ker}(\text{ad } \Lambda))$, is the set of $x \in \text{Ker}(\text{ad } \Lambda)$ such that $[x, y] = 0$ for any $y \in \text{Ker}(\text{ad } \Lambda)$, and, obviously, $\text{Cent}(\text{Ker}(\text{ad } \Lambda)) \subseteq \mathcal{H}[w]$. The two equivalent definitions of the flow on the right-hand-side of (2.11), together with the definition of Q as an intersection of two subspaces of $\widehat{g}$ characterized by the gradations $\mathbf{s}$ and $\mathbf{s}_w$, ensures that $\partial q(x)/\partial t_b^\theta \in Q$. In (2.11), $\text{P}_{\geq 0[\mathbf{s}_w]}$ and $\text{P}_{< 0[\mathbf{s}_w]}$ are the projectors onto $\widehat{g}_{\geq 0}(\mathbf{s}_w)$ and $\widehat{g}_{< 0}(\mathbf{s}_w)$, respectively.

Eqs.(2.11) define a unique flow on the gauge slice $Q^{(\text{can})}(x)$ because, then, the function $\theta(x)$ is fixed uniquely by the condition that $\partial L^{\text{can}}/\partial t_b \in Q^{(\text{can})}$, and $\theta(x)$ becomes a polynomial functional of the components of $q(x)$ and its derivatives. In (2.11), $V(x) \in \text{Im}(\text{ad } \Lambda) \cap \widehat{g}_{< 0}(\mathbf{s}_w)$ is the generator of the unique transformation which takes the Lax operator to the form

$$\mathcal{L} = \exp(\text{ad } V)(L) = L + [V, L] + \frac{1}{2}[V, [V, L]] + \dots = \partial_x + \Lambda + h(x), \quad (2.12)$$

² The dependence on x should be generalized to include all the t_b 's, but, as usual, we only show explicitly x .

where $h(x)$ is a function of x on $\text{Ker}(\text{ad } \Lambda) \cap \widehat{g}_{<i}(\mathbf{s}_w)$ (recall that i is the $\mathbf{s}_w$ -grade of Λ). It is important to remark that both $V(x)$ and $h(x)$ are polynomial functionals of the components of $q(x)$ and its x -derivatives³

One of the main properties of the flows defined by (2.11) is that they are Hamiltonian. Even more, they are bi-Hamiltonian when $\mathbf{s}$ is the homogeneous gradation [22]. The two Hamiltonian structures are defined on the set of gauge invariant functionals and they adopt the form of Kirillov-Poisson brackets on the affine Kac-Moody algebra $\widehat{g}$. If $\varphi[q]$ and $\psi[q]$ are two functionals of $q(x)$, the Poisson bracket defining the “first” Hamiltonian structure is

$$\{\varphi, \psi\}_1 = - \left(d_q \varphi, z^{-1} [d_q \psi, L] \right) , \quad (2.13)$$

while the bracket for the “second” structure is

$$\{\varphi, \psi\}_2 = (d_q \varphi_0, [d_q \psi_0, L]) - (d_q \varphi_{<0}, [d_q \psi_{<0}, L]) , \quad (2.14)$$

where the definitions of the inner product $(\cdot, \cdot)$ and of the functional derivative $d_q \varphi \in \widehat{g}$ are standard and can be found in [22]. Only those hierarchies for which $\mathbf{s} = \mathbf{s}_{\text{hom}} = (1, 0, \dots, 0)$ admit the two Hamiltonian structures, whereas the hierarchies are Hamiltonian only with respect to the second Poisson bracket in the general case. In these last two equations, we have used the following notation: superscripts denote $\mathbf{s}_w$ -grades, and subscripts indicate $\mathbf{s}$ -grades, but not just homogeneous grades as in [22]. For example, if $a \in \widehat{g}$ then a^j is the component of a on the subspace $\widehat{g}_j(\mathbf{s}_w)$, and a_k is its component on $\widehat{g}_k(\mathbf{s})$. We shall resort to this notation to simplify some expressions.

The Poisson brackets defined above admit non-trivial centres. In fact, notice that, by construction, the $\mathbf{s}$ -grade of the components of $q(x)$ is bounded, *i.e.*, there exists a non-negative integer n such that

$$\Lambda + q(x) \in \bigoplus_{j=0}^n \widehat{g}_j(\mathbf{s}) . \quad (2.15)$$

Then, it follows from (2.13) and (2.14) that the components of $q(x)$ whose $\mathbf{s}$ -grade is n are centres of the two Poisson bracket algebras; but they are not gauge invariant centres

³ In connection with the integrability of type II hierarchies, let us say that it should be possible to associate a flow not only to the elements of the centre of $\text{Ker}(\text{ad } \Lambda)$ but to any constant element of $\mathcal{H}[w]$ also in this case. The difference is that those flows not associated to the elements of the centre of $\text{Ker}(\text{ad } \Lambda)$ are always non-polynomial.

in general. In contrast, the existence of gauge invariant functionals that are centres of the Poisson brackets is directly related to the components of $h(x)$ in (2.12) with non-negative $\mathbf{s}_w$ -grade.

Proposition 1. *The functionals of the form $\Theta_f = (f(x), h(x))$ for $f(x)$ being a function of x on*

$$\text{Ker}(\text{ad } \Lambda)^{1-i} \cup \text{Cent} \left(\bigoplus_{j=1-i}^k \text{Ker}(\text{ad } \Lambda)^j \right) \quad (2.16)$$

are gauge invariant centres of the first Poisson bracket algebra for $k = 0$, and of the second Poisson bracket algebra for $k = -1$. $\square$

This result is the direct generalization of the Proposition 3.3 of [22] to include both type I and type II hierarchies; the proof is identical but taking into account that $\text{Ker}(\text{ad } \Lambda)$ is not abelian in the general case. Moreover, recall that the functionals Θ_f for $k = 0$ are non-dynamical and that $-x$ can be identified with the time t_Λ only when $P_{\geq 0[\mathbf{s}_w]}(h(x)) = 0$. The meaning of the mismatch between the centres of the two Poisson brackets has been clarified in [23].

The equations of the hierarchy are invariant under the scaling transformation

$$x \mapsto \lambda x, \quad t_{bj} \mapsto \lambda^{j/i} t_{bj}, \quad q^k \mapsto \lambda^{k/i-1} q^k, \quad (2.17)$$

where we have introduced a basis $\{e^k\}$ of Q , whose elements are graded with respect to $\mathbf{s}_w$, *i.e.*,

$$[d_{\mathbf{s}_w}, e^k] = k e^k, \quad \text{and} \quad q(x) = \sum_{k < i} q^k(x) e^k. \quad (2.18)$$

In relation to $\mathcal{W}$ -algebras, the crucial property is that (2.17) can be generalized to a conformal transformation

$$x \mapsto y(x), \quad q(x; z) \mapsto \tilde{q}(y(x); \tilde{z}), \quad (2.19)$$

with

$$q(x; z) = y' U[y] \left(\tilde{q}(y(x); \tilde{z}) \right) + \frac{1}{i} \left(\frac{y''}{y'} \right) H_{\mathbf{s}_w}, \quad (2.20)$$

and this transformation is a Poisson mapping of the second Poisson bracket. In (2.20),

$$y' = \frac{dy(x)}{dx}, \quad \tilde{z} = (y')^{-N_{\mathbf{s}_w}/i} z, \quad U[y] = (y')^{-\text{ad } H_{\mathbf{s}_w}/i},$$

and we have explicitly indicated the dependence of the potential on z , the affine parameter defining the loop algebra $\widehat{g}$. The infinitesimal version of (2.20) is $y(x) = x + \epsilon(x)$ and

$$\begin{aligned}
\delta_\epsilon q(x; z) &= q(x; z) - \widetilde{q}(x; z) \\
&= \partial_x (\epsilon(x) q(x; z)) - \frac{1}{i} \epsilon'(x) [N_{\mathbf{s}_w} z \frac{d}{dz} + H_{\mathbf{s}_w}, q(x; z)] + \frac{1}{i} \epsilon''(x) H_{\mathbf{s}_w} \\
&= [\partial_x + \Lambda(z) + q(x; z), \epsilon(x) (\Lambda(z) + q(x; z)) + \frac{1}{i} \epsilon'(x) H_{\mathbf{s}_w}] \\
&\quad - \frac{N_{\mathbf{s}_w}}{i} \epsilon'(x) [z \frac{d}{dz}, \Lambda(z) + q(x; z)],
\end{aligned} \tag{2.21}$$

which, using (2.18), is just

$$\begin{aligned}
\delta_\epsilon q(x; z) &= \sum_{k < i} e^k(z) \delta_\epsilon q^k(x) \\
&= \sum_{k < i} e^k(z) \left(\epsilon(x) \partial_x q^k(x) + \epsilon'(x) \left(1 - \frac{k}{i} \right) q^k(x) \right) + \frac{1}{i} \epsilon''(x) H_{\mathbf{s}_w}.
\end{aligned} \tag{2.22}$$

This transformation also induces a corresponding conformal transformation on the space of gauge equivalent classes, which is the phase space of the hierarchy.

3. The energy-momentum tensor.

The results of [22] ensure that the gauge invariant functionals of $q(x)$ close under the second Poisson bracket (2.14), and they have well defined properties with respect to the conformal transformation (2.20). This suggests a general relationship between the second Poisson bracket algebra and some (classical) quiral extended conformal algebras whose generators would be the components of $q^{(\text{can})}(x)$, the basis of gauge invariant functionals. This suggestion is supported by the examples analysed in [22]. A first step to establish the connection is to show that there is some gauge invariant functional $T_\epsilon[q]$ that plays the role of the energy-momentum tensor and generates the infinitesimal conformal transformation (2.21). However, let us anticipate that, when the second Hamiltonian structure is a $\mathcal{W}$ -algebra, it is defined by a conformal structure generally twisted with respect to the conformal structure associated to the transformation (2.21).

The condition that $T_\epsilon[q]$ generates (2.21) means that⁴

$$\begin{aligned} \delta_\epsilon q(x) &= \{T_\epsilon[q], q(x)\} \\ &= \left([\partial_x + \Lambda + q(x), (d_q T_\epsilon[q])_0] \right)_0 - \left([\Lambda + q(x), (d_q T_\epsilon[q])_{<0}] \right)_{>0}. \end{aligned} \quad (3.1)$$

The generator of the conformal transformation of the components of $q(x)$ whose $\mathbf{s}$ -grade is zero, $q_0(x)$, is easily derived from eqs.(2.21) and (3.1). The result is that

$$\delta_\epsilon q_0(x) = \{T_\epsilon[q], q(x)\}, \quad (3.2)$$

with

$$d_q T_\epsilon = \epsilon(x) (\Lambda + q(x))_0 + \frac{1}{i} \epsilon'(x) H_{\mathbf{s}_w}, \quad (3.3)$$

and the energy-momentum tensor is $T_\epsilon[q] = \int dx \epsilon(x) T(x)$ with

$$\begin{aligned} T(x) &= \frac{1}{2} \langle \Lambda + q(x), \Lambda + q(x) \rangle_{\hat{g}} - \frac{1}{i} \partial_x \langle H_{\mathbf{s}_w}, \Lambda + q(x) \rangle_{\hat{g}} \\ &= \frac{1}{2} \langle (\Lambda + q(x))_0, (\Lambda + q(x))_0 \rangle_{\hat{g}} - \frac{1}{i} \partial_x \langle H_{\mathbf{s}_w}, q_0(x) \rangle_{\hat{g}}, \end{aligned} \quad (3.4)$$

where $\langle \cdot, \cdot \rangle_{\hat{g}}$ is the Killing form of $\hat{g}$. In the particular case when $\mathbf{s}$ is the homogeneous gradation, this result has already been obtained in [23]. Notice that the components of $d_q T \in \hat{g}_{<0}(\mathbf{s})$ are not specified, and we have consistently taken them to zero. The Poisson bracket of two energy-momentum tensors can be calculated and the result is

$$\{T(x), T(y)\} = \frac{\langle H_{\mathbf{s}_w}, H_{\mathbf{s}_w} \rangle}{i^2} \delta'''(x-y) - 2 T(x) \delta'(x-y) - T'(x) \delta(x-y), \quad (3.5)$$

where $\delta^{(n)}(x-y) \equiv \partial_x^{(n)} \delta(x-y)$, and, now, $\langle \cdot, \cdot \rangle$ is the Killing form of the finite Lie algebra g . Eq.(3.5) shows that $T(x)$ indeed closes a classical Virasoro algebra with respect to the second Poisson bracket whose central extension is $12 \langle H_{\mathbf{s}_w}, H_{\mathbf{s}_w} \rangle / i^2$.

For the other components of $q(x)$, whose $\mathbf{s}$ -grade is > 0 , finding the generator of the conformal transformation from (2.21) and (3.1) is not so straightforward. In particular, we already know that some of these components are centres of the Poisson bracket algebra; so, there will be no energy-momentum tensor that generates their conformal transformation. As non-trivial examples involving $q_{>0}(x)$, let us mention the A_1 fractional KdV-hierarchies analysed in [22], and the integrable hierarchy associated to A_5 that we discuss in Section 7.

⁴ From now on we suppress the subscript indicating that the Poisson bracket is the second Poisson bracket, and the explicit indication of the dependence of Λ and q on z .

The reason why it is easy to derive the generator of the conformal transformation (3.2) is that the second Poisson bracket restricted to the functionals of $q_0(x)$ simplifies to

$$\{\varphi, \psi\} = \left([(d_q \varphi)_0, (d_q \psi)_0], (\Lambda + q(x))_0 \right) - \left((d_q \varphi)_0, \partial_x (d_q \psi)_0 \right), \quad (3.6)$$

which is just the Kirillov-Poisson bracket corresponding to the untwisted affinization in x of the finite Lie algebra $\widehat{g}_0(\mathbf{s})$, where the affine $\widehat{g}_0(\mathbf{s})$ current has been reduced to

$$J(x) = (\Lambda + q(x))_0; \quad (3.7)$$

as usual, the last term on the right-hand-side of (3.6) specifies the central extension. Notice that, because of lemma 1, either $(\Lambda)_0$ vanishes or it is a nilpotent element of $\widehat{g}_0(\mathbf{s})$. Of course, the energy-momentum tensor (3.4) comes from the Sugawara construction applied to the reduced current $J(x)$ improved by a term that depends on the gradation $\mathbf{s}_w$.

In the following, instead of considering the general case, we shall concentrate on the integrable hierarchies for which not only the condition (2.6) is satisfied, but also

$$\text{Ker}(\text{ad } (\Lambda)_0) \cap P = \emptyset, \quad (3.8)$$

which corresponds to the “non-degeneracy” condition of [10,11]; (3.8) is equivalent to $\text{Ker}(\text{ad } (\Lambda)_0) \cap \widehat{g}_{<0}(\mathbf{s}_w) = \emptyset$. Therefore, either P is empty or $(\Lambda)_0 \neq 0$. If $P = \emptyset$ then the hierarchy is a generalized mKdV hierarchy with $\mathbf{s} = \mathbf{s}_w$, $(\Lambda)_0 = 0$, and the second Poisson bracket algebra restricted to the functionals of $q_0(x)$ is an untwisted Kac-Moody algebra. In contrast, when $(\Lambda)_0 \neq 0$, we will show that this second Poisson bracket algebra is a $\mathcal{W}$ -algebra. In this case, a convenient choice of the gauge slice is:

Proposition 2. *Under the condition (3.8), the gauge slice $q^{(\text{can})}(x)$ can be chosen following the Drinfel’d-Sokolov prescription in such a way that*

$$\left(q^{(\text{can})>0}(x) \right)_0 = 0, \quad (3.9)$$

and that $q^{(\text{can})\leq 0}(x)$ depends only on $q_0(x)$.

Proof. First of all, notice that the proposition is obviously true when $(\Lambda)_0 = 0$ because $P = \emptyset$; then, (3.9) is an identity and $q^{(\text{can})\leq 0}(x) = q_0^{(\text{can})}(x) = q_0(x)$. Therefore, we shall

concentrate on the case of $(\Lambda)_0 \neq 0$. Consider the non-singular gauge transformation that takes $q(x)$ to the gauge slice $q^{(\text{can})}(x)$

$$q^{(\text{can})}(x) = \exp \left(\text{ad } \widehat{S} \right) \left(\partial_x + \Lambda + q(x) \right) - \partial_x - \Lambda , \quad (3.10)$$

with $\widehat{S}$ a polynomial functional of $q(x)$ and its x -derivatives. According to their $\mathbf{s}$ -grade, and taking into account that $\widehat{S} \in P \subset \widehat{g}_0(\mathbf{s})$, the components of (3.10) can be separated as

$$\left(q^{(\text{can})}(x) \right)_{>0} = \exp \left(\text{ad } \widehat{S} \right) \left((\Lambda + q(x))_{>0} \right) - (\Lambda)_{>0} , \quad (3.11)$$

$$\left(q^{(\text{can})}(x) \right)_0 = \exp \left(\text{ad } \widehat{S} \right) \left(\partial_x + (\Lambda + q(x))_0 \right) - \partial_x - (\Lambda)_0 . \quad (3.12)$$

Obviously $[(\Lambda)_0, P] \subset Q \cap \widehat{g}_0(\mathbf{s})$, and the condition (3.8) ensures that the map $\text{ad } (\Lambda)_0 : P \mapsto Q$ is injective. Consequently, it is possible to choose the gauge slice by applying the Drinfel'd-Sokolov prescription to (3.12). Within this choice, the generator $\widehat{S}$ of the gauge transformation (3.10) and the components of $(q^{(\text{can})}(x))_0$ depend only on $q_0(x)$, while those of $(q^{(\text{can})}(x))_{>0}$ depend on both $q_0(x)$ and $q_{>0}(x)$.

Now, let us analyse in detail the fact that $\text{ad } (\Lambda)_0 : P \mapsto Q_0$ is injective. This means that $\text{ad } (\Lambda)_0 : P^{-j} \mapsto Q_0^{i-j}$ is also injective for any positive integer j , which requires that

$$\dim \left(\widehat{g}_0^{-j} \right) \leq \dim \left(\widehat{g}_0^{i-j} \right) , \quad \text{for any } j \in \mathbb{Z} > 0 . \quad (3.13)$$

But $\dim \left(\widehat{g}_0^{-j} \right) = \dim \left(\widehat{g}_0^j \right)$, and it follows from (3.13) that

$$\dim \left(\widehat{g}_0^{-j} \right) = \dim \left(\widehat{g}_0^{i-j} \right) , \quad \text{for } 0 < j \in \mathbb{Z} < i . \quad (3.14)$$

We conclude that the number of components of $q(x) \in \widehat{g}_0(\mathbf{s}) \cap Q^{>0}$ equals the number of generators of gauge transformations in $P^{>-i}$, which, under (3.8), ensures the possibility of choosing the gauge slice such that (3.9) is satisfied. This completes the proof. $\square$

It is well known that the construction of $\mathcal{W}$ -algebras is based on the $sl(2, \mathbb{C})$ subalgebras of a finite Lie algebra [8], while the integrable hierarchies of [21,22] are constructed in terms of the different Heisenberg subalgebras of a loop algebra. The form of the reduced current (3.7) suggests the kind of connection between the two constructions that should be investigated.

If $(\Lambda)_0$ does not vanish, it is a nilpotent element of $\widehat{g}$. Then the Jacobson-Morozov theorem —theorem 17 of [32], pag. 100— affirms that there exists a unique element $J_- \in \widehat{g}_0(\mathbf{s}) \cap \widehat{g}_{-i}(\mathbf{s}_w)$ such that $J_+ = (\Lambda)_0$, J_- , and $J_0 = [J_+, J_-]$ close an $A_1 = sl(2, \mathbb{C})$ subalgebra of $\widehat{g}_0(\mathbf{s})$. J_0 and the $H_{\mathbf{s}}$ that defines (2.7) lie on the same Cartan subalgebra of g . Their relation is summarized by the following results.

Lemma 2. ([33]) *Let $S = (J_+, J_0, J_-)$ be the A_1 subalgebra of $\widehat{g}_0(\mathbf{s})$ with $J_+ = (\Lambda)_0$. Then, $H_{\mathbf{s}_w} = J_0 + Y$, and the generator Y commutes with all the generators of S . $\square$*

Hence, if Y does not vanish, $J_+ = (\Lambda)_0$, $H_{\mathbf{s}_w}$, and J_- generate an $A_1 \oplus u(1)$ subalgebra of $\widehat{g}_0(\mathbf{s})$. Under the adjoint action of this subalgebra, $\widehat{g}_0(\mathbf{s})$, considered as a vector space, decomposes as the direct sum of a finite number of irreducible representations of A_1

$$\widehat{g}_0(\mathbf{s}) = \bigoplus_{k=1}^n D_{j_k}(y_k), \quad (3.15)$$

where j_k and y_k are the spin and the eigenvalue of Y on the irreducible representation $D_{j_k}(y_k)$. Notice that the $\mathbf{s}_w$ -grade of Λ is i , thus, the normalization of the A_1 subalgebra used in lemma 2 is

$$[J_0, J_{\pm}] = \pm i J_{\pm}, \quad [J_+, J_-] = J_0, \quad (3.16)$$

and the eigenvalues of J_0 on $D_{j_k}(y_k)$ are $(-ij_k, -i(j_k - 1), \dots, +ij_k)$, while those of $H_{\mathbf{s}_w}$ are $(y_k - ij_k, y_k - i(j_k - 1), \dots, y_k + ij_k)$.

Lemma 3. ([33]) *Under the condition (3.8), the eigenvalues of Y in the decomposition (3.15) are bounded according to*

$$|y_k| \leq i j_k. \quad \square \quad (3.17)$$

Proposition 2 ensures that the components of $q^{(\text{can}) \leq 0}(x)$ depend only on $q_0(x)$, but it only specifies the components of the gauge slice whose $\mathbf{s}_w$ -grade is > 0 ; they correspond to the gauge fixing of the transformations generated by $P^{>-i}$. The residual gauge freedom is generated by the elements of $P^{\leq -i}$, which just induces the transformation of $q^{\leq 0}(x)$ according to (3.12). If (3.8) is fulfilled, that residual freedom can be fixed using the Drinfel'd-Sokolov prescription in such a way that $q^{(\text{can}) \leq 0}(x)$ is a combination of the lowest

weights in the decomposition (3.15); *i.e.*, of those elements of $Q^{\leq 0}$ that cannot be expressed as $[J_+, *]$, $J_+ = (\Lambda)_0$. All this proves the following.

Theorem 1. *Under the condition (3.8), the gauge slice can be chosen as*

$$q^{(\text{can})}(x) = q_{>0}^{(\text{can})}(x) + \mathcal{J}(x), \quad (3.18)$$

where $\mathcal{J}(x)$ is a combination of all the lowest weights of the irreducible representations of A_1 into the decomposition (3.15) of $\widehat{g}_0(\mathbf{s})$. The components of $\mathcal{J}(x)$ are polynomial functionals of $q_0(x)$ and their x -derivatives, while those of $(q^{(\text{can})}(x))_{>0}$ depend also on $q_{>0}(x)$. $\square$

Remark. The form of $\mathcal{J}(x)$ depends only on $(\Lambda)_0$ because $J_+ = (\Lambda)_0$ fixes the A_1 subalgebra of $\widehat{g}_0(\mathbf{s})$ inducing the decomposition (3.15), and lemma 3 ensures that all the lowest weights of the representations of A_1 indeed have not positive $\mathbf{s}_w$ -grade. Thus, as far as (3.8) is satisfied, the form of $(q^{(\text{can})}(x))_0$ is independent of the choice of $\mathbf{s}_w$.

Let us go back to the form of the energy-momentum tensor. We have already said that (3.4) comes from the Sugawara construction applied to the reduced current (3.7), improved by a term that depends on the gradation $\mathbf{s}_w$, and that it generates the conformal transformation (2.19) on $q_0(x)$. Nevertheless, in general, this energy-momentum tensor is not gauge invariant. Consider the gauge transformation

$$q(x) \mapsto \widetilde{q}(x) = \exp(\text{ad } S)(\partial_x + \Lambda + q(x)) - \partial_x - \Lambda,$$

then

$$\begin{aligned} T_\epsilon[\widetilde{q}] &= \frac{1}{2} \int dx \, \epsilon(x) \langle \exp(\text{ad } S)(\partial_x + \Lambda + q(x)) - \partial_x, \exp(\text{ad } S)(\partial_x \\ &\quad + \Lambda + q(x)) - \partial_x \rangle_{\widehat{g}} \\ &\quad + \frac{1}{i} \int dx \, \epsilon'(x) \langle H_{\mathbf{s}_w}, \exp(\text{ad } S)(\partial_x + \Lambda + q(x)) - \partial_x \rangle_{\widehat{g}} \\ &= T_\epsilon[q] - \int dx \, \epsilon(x) \langle \exp(-\text{ad } S)(\partial_x) - \partial_x, \Lambda + q(x) \rangle_{\widehat{g}} \\ &\quad + \frac{1}{i} \int dx \, \epsilon'(x) \langle \exp(-\text{ad } S)(H_{\mathbf{s}_w}) - H_{\mathbf{s}_w}, \Lambda + q(x) \rangle_{\widehat{g}} \\ &\equiv T_\epsilon[q] + \Sigma_\epsilon(S; q), \end{aligned} \quad (3.19)$$

where we have used the invariance of the bilinear form of $\widehat{g}$. Now, using (2.2) and (2.4), it can be checked that $\Sigma_\epsilon(S; q)$ depends only on the components of $S \in P^{>-i}$ and on those of $q \in Q^{>0}$. Therefore, $\Sigma_\epsilon(S; q)$ vanishes when $i = 1$ and the energy-momentum tensor (3.4) is gauge invariant only in this case.

In contrast, when $i > 1$ the energy-momentum tensor (3.4) is not gauge invariant. Recall that (3.4) has been derived from the condition that it generates the conformal transformation of $q_0(x)$. As we only need the generator of the conformal transformation of the gauge invariant functionals of $q_0(x)$, $T_\epsilon[q]$ can be changed for any other functional $\widehat{T}_\epsilon[q]$ such that its functional derivative equals (3.3) up to the generator of an infinitesimal gauge transformation; *i.e.*,

$$(d_q \widehat{T}_\epsilon)_0 - (d_q T_\epsilon)_0 \in P. \quad (3.20)$$

The natural choice is:

Proposition 3. *For any choice of the gauge slice $Q^{(\text{can})}$, the gauge invariant functional*

$$\Theta_\epsilon[q] = \int dx \, \epsilon(x) \, \Theta(x) = T_\epsilon[q^{(\text{can})}], \quad (3.21)$$

generates the conformal transformation of the gauge invariant functionals of $q_0(x)$.

Proof. To start with, let us recall that the components of $q^{(\text{can})}(x)$ are understood as gauge invariant functionals of $q(x)$, and that we fix the gauge following the Drinfel'd-Sokolov prescription. The relation between the components of $q^{(\text{can})}(x)$ whose $\mathfrak{s}_w$ -grade is j and $q(x)$ is

$$\begin{aligned} q^{(\text{can})j}(x) &= \left(\exp(\text{ad } \widehat{S}) (\partial_x + \Lambda + q(x)) - \partial_x - \Lambda \right)^j \\ &= q^j(x) - [\Lambda, \widehat{S}^{j-i}(x)] + \Delta(\widehat{S}^{>j-i}; q^{>j}). \end{aligned} \quad (3.22)$$

The components of $\widehat{S}$ are functionals of $q(x)$ fixed by the condition that $q^{(\text{can})}(x) \in Q^{(\text{can})}$, with $Q^{(\text{can})}$ some complementary vector space of $[\Lambda, P]$ in Q ; so, $\widehat{S}^{-j}(x)$ depends only on the components of $q(x)$ whose $\mathfrak{s}_w$ -grade is $\geq i - j$. Moreover, (3.22) also shows that $q^{(\text{can})j}(x)$ is a functional only of the components of $q(x)$ with $\mathfrak{s}_w$ -grade $\geq j$. Now, let us consider (3.19) with $S = \widehat{S}$; since $\Sigma_\epsilon(\widehat{S}; q)$ depends only on the components of $q(x)$ whose $\mathfrak{s}_w$ -grade is positive, we conclude that $(d_q \Sigma_\epsilon(\widehat{S}; q))_0 \in P$, which, using (3.20), proves the proposition. $\square$

Finally, with the choice of the gauge slice of theorem 1, the gauge invariant energy-momentum tensor (3.21) becomes

$$\begin{aligned}\Theta(x) &= \frac{1}{2} \langle \Lambda + q^{(\text{can})}(x), \Lambda + q^{(\text{can})}(x) \rangle_{\hat{g}} - \frac{1}{i} \partial_x \langle H_{\mathbf{s}_w}, \Lambda + q^{(\text{can})}(x) \rangle_{\hat{g}} \\ &= \langle (\Lambda)_0, \mathcal{J}^{-i}(x) \rangle_{\hat{g}} + \frac{1}{2} \langle \mathcal{J}^0(x), \mathcal{J}^0(x) \rangle_{\hat{g}} - \frac{1}{i} \partial_x \langle H_{\mathbf{s}_w}, \mathcal{J}^0(x) \rangle_{\hat{g}},\end{aligned}\tag{3.23}$$

which shows that $\Theta(x)$ is a gauge invariant functional of $q_0(x)$.

4. The second Poisson bracket and the $\mathcal{W}$ -algebras.

Theorem 1 shows that, under condition (3.8), the components of $\mathcal{J}(x)$ provide a basis for the gauge invariant functionals of the components of $q(x)$ whose $\mathbf{s}$ -grade is zero. Obviously, the second Poisson bracket of two of these functionals is another gauge invariant functional of $q_0(x)$ and it makes sense to consider the restriction of the second Poisson bracket algebra to these functionals. Recall that, in the previous section, the components of $q^{(\text{can})}(x)$ have to be understood as functionals of $q(x)$, hence it would be more precise to write eq.(3.18) as

$$q^{(\text{can})}\left(q(x)\right) = q_{>0}^{(\text{can})}\left(q_0(x), q_{>0}(x)\right) + \mathcal{J}\left(q_0(x)\right),\tag{4.1}$$

and it follows from the gauge fixing prescription that $q^{(\text{can})}(q(x)) = q(x)$ if $q(x) \in Q^{(\text{can})}$; in particular, $\mathcal{J}(q_0(x)) = q_0(x)$ if $q_0(x) \in Q_0^{(\text{can})}$.

In the previous section, we have proved that the restriction of the second Poisson bracket algebra to the gauge invariant functionals of $q_0(x)$ is just the Hamiltonian reduction of the untwisted affinization of $\hat{\mathfrak{g}}_0(\mathbf{s})$ defined by the bracket (3.6), and corresponding to

$$J(x) = (\Lambda)_0 + \mathcal{J}(x),\tag{4.2}$$

which is one of the $\mathcal{W}$ -algebras considered in [8,10,11]. In particular, it is the $\mathcal{W}$ -algebra associated to the A_1 subalgebra of $\hat{\mathfrak{g}}_0(\mathbf{s})$ whose nilpotent J_+ element is $(\Lambda)_0$.

Notice that the relevant conformal structure in terms of which all the generators of the $\mathcal{W}$ -algebra are primary fields is defined by the semisimple element of the A_1 subalgebra J_0 , and the appropriated energy-momentum tensor is

$$\mathcal{T}(x) = \langle (\Lambda)_0, \mathcal{J}^{-i}(x) \rangle_{\hat{g}} + \frac{1}{2} \langle \mathcal{J}^0(x), \mathcal{J}^0(x) \rangle_{\hat{g}},\tag{4.3}$$

such that [10]

$$\{\mathcal{T}_\epsilon[q], \mathcal{J}(x)\} = \epsilon(x)\mathcal{J}'(x) + \epsilon'(x) \left(\mathcal{J}(x) - \frac{1}{i} [J_0, \mathcal{J}(x)] \right) - \frac{1}{i} \epsilon'''(x) J_- , \quad (4.4)$$

where we use the normalization (3.16), and $\mathcal{T}_\epsilon[q] = \int dx \epsilon(x) \mathcal{T}(x)$. $\mathcal{T}(x)$ closes a classical Virasoro algebra whose central extension is $12\langle J_+, J_- \rangle / i = 12\langle J_0, J_0 \rangle / i^2$. As the elements of $\mathcal{J}(x)$ are the lowest weights in (3.15), $\langle J_0, \mathcal{J}(x) \rangle_{\hat{g}} = 0$ and $\mathcal{T}(x)$ equals $\Theta(x)$ only if $H_{\mathbf{s}_w}$ is proportional to J_0 ; otherwise,

$$\mathcal{T}(x) - \Theta(x) = \frac{1}{i} \partial_x \langle H_{\mathbf{s}_w} - J_0, \mathcal{J}^0(x) \rangle_{\hat{g}} . \quad (4.5)$$

$\Theta(x)$, given by (3.23), defines the conformal structure corresponding to the transformation (2.20) that generalizes the scaling transformation (2.17) under which the integrable hierarchy is homogeneous. Hence, in general, the conformal structure of the $\mathcal{W}$ -algebra is twisted with respect to the conformal structure defining the scaling properties of the integrable hierarchy.

In relation to this, it is convenient to consider again the remark below theorem 1. It indicates that, as long as (3.8) keeps up, the second Poisson bracket algebra does not depend on the choice of the gradation $\mathbf{s}_w$. We have anticipated in Section 2 that, for a given Heisenberg subalgebra $\mathcal{H}[w]$, the choice of $\mathbf{s}_w$ is not unique, a question that has not been mentioned at all in [21,22]. The origin of this ambiguity can be easily understood by noticing that, for a given choice of $\mathbf{s}_w$, the elements of $\mathcal{H}[w]$ are also graded by any other gradation $\mathbf{s}'$ such that

$$\frac{H_{\mathbf{s}_w}}{N_{\mathbf{s}_w}} - \frac{H_{\mathbf{s}'}}{N_{\mathbf{s}'}} \in \mathcal{H}[w] . \quad (4.6)$$

In fact, it can be proven that these are the only ambiguities in the choice of $\mathbf{s}_w$, and that they are obviously linked to the existence of elements of $\mathcal{H}[w]$ whose $\mathbf{s}_w$ -grade is zero. Moreover, notice that (4.6) ensures that the grade of the elements of $\mathcal{H}[w]$ is uniquely defined up to multiplication by a global rational factor, in agreement with lemma 1. This shows, in particular, that the scaling dimensions of the times t_{b_j} in (2.17) are also uniquely defined. Let us consider the Heisenberg subalgebras of $\hat{g}$ as the affinization in z of a fixed Cartan subalgebra of g [31]. The different Heisenberg subalgebras arise in relation to the different finite order (inner) automorphisms of g , which are classified by the conjugacy classes of the Weyl group of g [31]. The connection between a Weyl group element, say w

up to conjugacy, and the associated Heisenberg subalgebra $\mathcal{H}[w]$ is that there is a lift of w onto $\widehat{g}$, denoted $\widehat{w}$, which acts on $\mathcal{H}[w]$ as

$$\widehat{w}(b_j) \equiv w(b_j) = \exp\left(\frac{2\pi i}{N} j\right) b_j, \quad (4.7)$$

with N being the order of w , $w^N = 1$. The different choices of $\mathbf{s}_w$ correspond precisely to the different choices of $\widehat{w}$, which, in the “shifted picture”, can be expressed as [34]

$$\widehat{w} = \exp\left(\frac{2\pi i}{N_{\mathbf{s}_w}} \text{ad } H_{\mathbf{s}_w}\right). \quad (4.8)$$

The choice of $\mathbf{s}_w$ is unique for the principal Heisenberg subalgebra, which is related to the conjugacy class of the Coxeter element of the Weyl group of g . In contrast, the choice of $\mathbf{s}_w$ for the homogeneous Heisenberg subalgebra, for which w is the identity, is completely arbitrary. In [31], a canonical choice of $\mathbf{s}_w$ is determined by

$$\langle H_{\mathbf{s}_w}, b_j \rangle = 0, \quad \text{for any } b_j \in \mathcal{H}[w]; \quad (4.9)$$

this condition is non-trivial only for $b_j \in \mathcal{H}[w] \cap \widehat{g}_0(\mathbf{s}_w) \subset \mathcal{H}[w] \cap \widehat{g}_0(\mathbf{s})$. Let us assume that $H_{\mathbf{s}_w} \in \text{Im}(\text{ad } (\Lambda)_0)$, *i.e.*, that there exists some $M \in \widehat{g}_0(\mathbf{s}) \cap \widehat{g}_{-i}(\mathbf{s}_w)$ such that $H_{\mathbf{s}_w} = [(\Lambda)_0, M]$. Then, the invariance of the bilinear form of $\widehat{g}$ implies that $\langle H_{\mathbf{s}_w}, b_j \rangle = \langle [(\Lambda)_0, M], b_j \rangle$ vanishes and (4.9) can be viewed as a necessary condition to ensure that $H_{\mathbf{s}_w}$ is proportional to J_0 .

All these results fix the connection between the algebras defined by the second Poisson bracket of the integrable hierarchies of [21,22] and the $\mathcal{W}$ -algebras of [8,10,11].

Theorem 2. *Let us consider the integrable hierarchy associated to the data*

$$\{\widehat{g}, \mathcal{H}[w], s_w, \mathbf{s}, \Lambda\}, \quad \text{such that } \text{Ker}(\text{ad } (\Lambda)_0) \cap P = \emptyset.$$

(i) If $(\Lambda)_0 \neq 0$, the restriction of the second Poisson bracket algebra to the gauge invariant functionals of $q_0(x)$ is the $\mathcal{W}$ -algebra associated to the A_1 subalgebra of $\widehat{g}_0(\mathbf{s})$ specified by the nilpotent element $J_+ = (\Lambda)_0$. (ii) Otherwise, if $(\Lambda)_0 = 0$, such restriction is just the Kac-Moody algebra corresponding to the untwisted affinization of $\widehat{g}_0(\mathbf{s})$. $\square$

Together with this theorem, let us point out, first, that theorem 2 does not specify the Poisson brackets involving the components of $q_{>0}^{(\text{can})}(x)$; in any case, proposition 1 shows

that some of these components are centres of the Poisson bracket algebra. Second, in general, the gradation $\mathbf{s}_w$ cannot be chosen such that $H_{\mathbf{s}_w}$ is proportional to J_0 ; therefore, the conformal transformations generated by $\mathcal{T}(x)$ will be different from the transformation induced by (2.19) and (2.20). Anyway, it follows from lemma 2 that the condition (3.8) is always satisfied in those particular cases when $H_{\mathbf{s}_w}$ is proportional to J_0 . So, it is interesting to consider the following straightforward result.

Lemma 4. *Let us suppose that $H_{\mathbf{s}_w}$ is proportional to J_0 . Then, either $i = 1$ or $i = 2$, and the value $i = 1$ means that the corresponding embedding of A_1 into g is integral.*

Proof. Let us look at the decomposition (3.15). According to lemma 3, and with the normalization used there, $H_{\mathbf{s}_w}$ being proportional to J_0 means that, actually, $H_{\mathbf{s}_w} = J_0$, or, equivalently, that $Y = 0$. Then, the gradation induced by $\mathbf{s}_w$ is

$$\widehat{g} = \bigoplus_{k \in I} \widehat{g}_k(\mathbf{s}_w),$$

where

$$I = \bigcup_{k=1}^n \{-ij_k, -i(j_k - 1), \dots, i(j_k - 1), ij_k\} + \mathbb{Z} N_{\mathbf{s}_w},$$

which means that all the grades are proportional to i . Since $\mathbf{s}_w$ is an integer gradation of $\widehat{g}$, and the non-zero grades are relatively prime, there are only two possibilities. The first one is that all the representations in (3.15) have integer spin; in this case $i = 1$. The second possibility is that there are representations whose spin is half-integer, then $i = 2$. $\square$

Theorem 2 generalizes the well known relationship between the Gel'fand-Dickey algebras and the $\mathcal{W}$ -algebras [5,7] that we have summarized in the introduction. The logical question is if all the $\mathcal{W}$ -algebras that can be constructed out of the inequivalent A_1 subalgebras can be recovered in this way. This requires that, for any A_1 subalgebra of $\widehat{g}_0(\mathbf{s})$, J_+ can be expressed as the zero $\mathbf{s}$ -grade projection of a $\mathbf{s}_w$ -graded element of some Heisenberg subalgebra of $\widehat{g}$. Hence, it is necessary to investigate in detail the relation between the embeddings of A_1 into a Lie algebra g and the conjugacy classes of the Weyl group of g that classify the inequivalent Heisenberg subalgebras of $g^{(1)}$. A formal relation of this kind has been realized by mathematicians [35] after the classification of the conjugacy classes of the Weyl group was completed, and, more recently, this topic has attracted the

interest of physicists in connection with the structure of integrable two-dimensional field theories [29,30].

Before addressing this question in the next section, let us settle the two following results that will be useful when $\mathbf{s}$ is the homogeneous gradation.

Lemma 5. *The components of $\hat{g}$ whose homogeneous grade is $\geq m$ have $\mathbf{s}_w$ -grade $\geq (\mathbf{s}_w)_0 + (m-1)N_{\mathbf{s}_w}$.*

Proof. Notice that the $\mathbf{s}_w$ -grade of any element of g ($\equiv g \otimes z^0$) is $\geq$ than the grade of the lowest root's step operator, *i.e.*, $\geq$ than

$$-\sum_{j=1}^r k_j(\mathbf{s}_w)_j = (\mathbf{s}_w)_0 - N_{\mathbf{s}_w} , \quad (4.10)$$

where we have used that the Kac label $k_0 = 1$ for the untwisted Kac-Moody algebras. The result follows just by considering the $\mathbf{s}_w$ -grade of the elements $a \otimes z^n \in \hat{g}$ with $n \geq m$. $\square$

Lemma 6. *The minimal positive $\mathbf{s}_w$ -grade of the elements of $\mathcal{H}[w]$ is $\geq (\mathbf{s}_w)_0$. Moreover, if $\mathbf{s}_w$ is $\succeq$ than the homogeneous gradation, then $(\mathbf{s}_w)_0 \geq 1$.*

Proof. To define the gradations, we have chosen a particular Cartan subalgebra H of g in such a way that the elements of H have zero grade with respect to any gradation (see (2.7)). Besides, if $B_{\pm}$ are the maximal positive and negative nilpotent subalgebras of g with respect to H , *i.e.*, $g = B_+ \oplus H \oplus B_-$, the elements of B_+ and B_- have, respectively, grade ≥ 0 and ≤ 0 with respect to whatever gradation. Recall now that the elements of $\mathcal{H}[w]$ are semisimple while those of $B_{\pm}$ are nilpotent. Hence, any element of $\mathcal{H}[w]$ with positive $\mathbf{s}_w$ -grade has to be of the form $z^n \otimes \alpha_+ + z^{n+1} \otimes \alpha_-$, with $n \geq 0$, and $\alpha_{\pm} \in B_{\pm} \neq 0$. Therefore, the first part of the lemma follows by using lemma 5 with $m = 1$, and if $\mathbf{s}_w$ is $\succeq$ than the homogeneous gradation, then $(\mathbf{s}_w)_0 \neq 0$, *i.e.*, $(\mathbf{s}_w)_0 \geq 1$. $\square$

These two lemmas lead to

Lemma 7. *When $\mathbf{s}$ is the homogeneous gradation, $(\Lambda)_0 \neq 0$ if and only if*

$$(\mathbf{s}_w)_0 \leq i < N_{\mathbf{s}_w} . \quad (4.11)$$

Moreover, if $i = (\mathbf{s}_w)_0$, the $\mathbf{s}$ -grade of all the components of the potential $q(x)$ is zero, i.e., $q(x) \in \widehat{g}_0(\mathbf{s})$, and the two conditions (2.6) and (3.8) are equivalent. $\square$

As a remark, let us mention that, in general, not all Heisenberg subalgebras have generators whose $\mathbf{s}_w$ -grade equals the lower bound $i = (\mathbf{s}_w)_0$.

5. Embeddings of A_1 and conjugacy classes of the Weyl group.

In this section, we restrict ourselves to the A_n algebras. Similar results for the other classical Lie algebras will be presented elsewhere.

5.1 The A_1 subalgebras of A_n .

The embeddings of A_1 into a semisimple Lie algebra have been classified originally by E.B. Dynkin [36], a classification that has been made explicit for those algebras of rank up to 6 in [37] —see also [11,38] for recent reviews of this classification inspired by the construction of $\mathcal{W}$ -algebras and non-abelian Toda models. Here, we normalize the A_1 subalgebras as

$$[J_0, J_{\pm}] = \pm 2 J_{\pm}, \quad [J_+, J_-] = J_0. \quad (5.1)$$

The different embeddings of A_1 into the Lie algebra g correspond to the different injective homomorphisms $\iota : A_1 \mapsto g$, and, in practice, we shall identify A_1 with its image under ι . An embedding of A_1 is specified by its defining vector $\mathbf{f} = (f_1, \dots, f_r)$, which provides the decomposition of J_0 with respect to a basis of the Cartan subalgebra $\{H_1, \dots, H_r\}$ of g ,

$$J_0 = \sum_{j=1}^r f_j H_j; \quad (5.2)$$

obviously, r is the rank of the Lie algebra g . It can be proved that two embeddings of A_1 are conjugated if and only if their J_0 's are conjugated under some automorphism of g [36]. Even more, through conjugation with the Weyl group of g , the basis of simple roots $\{\alpha_1, \dots, \alpha_r\}$ can be chosen such that the quantities $c_j = \alpha_j(J_0) \geq 0$; in fact, with the normalization (5.1), these numbers can only take the values 0, 1, or 2. Considering the decomposition of the Lie algebra g , as a vector space, under the adjoint action of $\iota(A_1)$, the value of

$c_j = 2, 1$, or 0 indicates that the step operator E_{α_j} belongs to a representation whose spin is integer, half-integer, or zero, respectively. The numbers $c_1, \dots, c_r$ are associated to the nodes of the Dynkin diagram of g , and this set of numbers is called the “characteristic” of the embedding.

A particularly important example is provided by the “principal embedding” defined as [28]

$$J_+^{(\text{p})} = \sum_{j=1}^r \lambda_j E_{\alpha_j}, \quad J_0^{(\text{p})} = \sum_{j=1}^r \frac{4\omega_j}{\alpha_j^2} \cdot H, \quad (5.3)$$

where the ω_j ’s are the fundamental weights of g defined as $\alpha_j \cdot \omega_k = (\alpha_j^2/2)\delta_{jk}$. The λ_j ’s are non-vanishing arbitrary constants, usually taken to be 1, but notice that J_0 is completely independent of them. The “characteristic” of the principal embedding is $\{2, 2, \dots, 2\}$.

For A_n , it is convenient to represent the Cartan subalgebra using an $(n+1)$ -dimensional Euclidean space as follows [11]. Let $\{e_1, \dots, e_{n+1}\}$ be a basis of this space that is orthonormal with respect to the Cartan-Killing form. In this basis, the roots of A_n are $(e_i - e_j)$, where $i, j = 1, \dots, n+1$, and a suitable choice of the simple roots is $\{\alpha_j = e_j - e_{j+1}, j = 1, \dots, n\}$. Accordingly, the basis of the Cartan subalgebra can be chosen in terms of the set $\{H_1, \dots, H_{n+1}\}$ such that

$$[H_i, E_{(e_j - e_k)}] = (\delta_{i,j} - \delta_{i,k})E_{(e_j - e_k)}, \quad i, j, k = 1, \dots, n+1, \quad (5.4)$$

and the elements of the Cartan subalgebra are $\sum_{j=1}^{n+1} p_j H_j$ with $\sum_{j=1}^{n+1} p_j = 0$. Having made these choices, the defining vector of the principal embedding of A_1 into A_n is

$$\mathbf{f}^{\text{p}} = (n, n-2, \dots, 2-n, -n). \quad (5.5)$$

Dynkin’s result is that all the non-conjugated embeddings of A_1 into A_n can be described as the principal A_1 embedding of a regular subalgebra⁵ of A_n . Dynkin has also classified the regular subalgebras of semisimple Lie algebras [36]. In the case of A_n , they are in one-to-one relation with the partitions of $n+1$, and the regular subalgebra of A_n associated to

$$n+1 = \sum_{j=1}^k n_j, \quad n_k \geq n_{k-1} \geq \dots \geq n_1, \quad (5.6)$$

⁵ Consider the root space decomposition of a Lie algebra g with respect to a Cartan subalgebra H : $g = H \oplus_{\alpha \in \Delta} g_{\alpha}$, where Δ is the root system of g . Then, a subalgebra $g' \subset g$ is called regular if it can be seen, up to conjugation, as $g' = H' \oplus_{\alpha \in \Delta'} g_{\alpha}$, with $\Delta' \subset \Delta$ and $H' \subset H$.

where $k \geq 1$ and n_j are positive integer numbers, is

$$\bigoplus_{\substack{j=1 \\ n_j \neq 1}}^k A_{n_j-1} \subset A_n . \quad (5.7)$$

Then, the corresponding embedding of A_1 into A_n is defined by

$$J_{\pm}^{(n_k, \dots, n_1)} = \sum_{\substack{j=1 \\ n_j \neq 1}}^k \left(J_{\pm}^{(p)} \right)_{A_{n_j-1}} , \quad J_0^{(n_k, \dots, n_1)} = \sum_{\substack{j=1 \\ n_j \neq 1}}^k \left(J_0^{(p)} \right)_{A_{n_j-1}} . \quad (5.8)$$

The defining vector of this embedding is just the union of the defining vectors of the principal embeddings of each A_{n_j-1} factor in (5.7) plus the required number of zeroes to complete the $n+1$ components of $\mathbf{f}$ —the number of zeroes equals the number of 1's in (5.6). The components of the resulting $\mathbf{f}$ can be arranged such that $f_{n+1} \leq f_n \leq \dots \leq f_1$, which corresponds to the change of the basis of simple roots leading to the “characteristic” of the embedding [11]. Finally, after the components of $\mathbf{f}$ have been ordered, the components of the “characteristic” are $c_j = f_j - f_{j+1}$, $j = 1, \dots, n$.

For instance, the defining vector of the embedding associated to the partition $4 = 2 + 1 + 1$, *i.e.*, to the regular subalgebra $A_1 \subset A_3$, is $(1, 0, 0, -1)$. Another example, the defining vector corresponding to $7 = 4 + 2 + 1$, *i.e.*, to $A_3 \oplus A_1 \subset A_6$, is $(3, 1, 1, 0, -1, -1, -3)$, as follows from the union of $(3, 1, -1, -3)$, $(1, -1)$, and (0) . In these two examples, the “characteristics” are $(1, 0, 1)$ and $(2, 0, 1, 1, 0, 2)$, respectively.

Notice that, for any embedding of A_1 into A_n , the components of $\mathbf{f}$ satisfy the symmetry relation $f_j = -f_{n+2-j}$, which implies that the components of the “characteristic” satisfy $c_j = c_{n+1-j}$.

5.2 The conjugacy classes of the Weyl group of A_n and the Heisenberg subalgebras of $A_n^{(1)}$.

It has been shown by V.G. Kac and D.H. Peterson that the inequivalent Heisenberg subalgebras of the affine untwisted Kac-Moody algebra $g^{(1)}$ are in one-to-one relation with the conjugacy classes of the Weyl group of g [31]—we have briefly summarized the connection between the elements of the Weyl group of g and the Heisenberg subalgebras of $g^{(1)}$ in (4.7) and (4.8). The complete classification of the conjugacy classes of the Weyl group, including classical and exceptional Lie algebras, has been developed by R.W. Carter [39].

In connection with the construction of the twisted vertex operator representations of Kac-Moody algebras, more recently, very detailed descriptions of the conjugacy classes of the Weyl group and the Heisenberg subalgebras of the untwisted affinizations of the classical Lie algebras have been presented in [40, 41] (see also [19, 25]).

The Weyl group of A_n is isomorphic to S_{n+1} , the group of permutations of $n + 1$ elements. Therefore, it can be represented as a set of linear transformations acting on an $(n + 1)$ -dimensional Euclidean space with an orthonormal basis $\{e_1, \dots, e_{n+1}\}$ —the same we have used to describe the Cartan subalgebra of A_n . The elements of the Weyl group of A_n are permutations of the elements of this basis; it can be shown that the conjugacy classes of the Weyl group of A_n correspond to permutations without specifying the particular sets of e_j 's on which they act. The conjugacy classes of the Weyl group of A_n are too in one-to-one relation with the partitions of $n + 1$. The conjugacy class corresponding to the partition (5.6) is usually denoted as $[n_1, n_2, \dots, n_k]$, where the order of the n_j 's is irrelevant. If the elements of the basis are divided into k subsets with $n_1, n_2, \dots, n_k$ elements, respectively, the conjugacy class consists in a cyclic permutation in each of the subsets, *i.e.*,

$$\begin{aligned}
& e_1 \rightarrow e_2 \rightarrow \dots \rightarrow e_{n_1} \rightarrow e_1 \\
& e_{n_1+1} \rightarrow e_{n_1+2} \rightarrow \dots \rightarrow e_{n_1+n_2} \rightarrow e_{n_1+1} \\
& \dots\dots\dots \\
& e_{n_1+\dots+n_{k-1}+1} \rightarrow e_{n_1+\dots+n_{k-1}+2} \rightarrow \dots \rightarrow e_{n+1} \rightarrow e_{n_1+\dots+n_{k-1}+1} .
\end{aligned} \tag{5.9}$$

A particularly important element of the Weyl group is the Coxeter one, and the conjugacy class of the Coxeter element of A_n is just $[n + 1]$. It can be shown that the eigenvalues of any element of $[n + 1]$ are $\{\omega^j, j = 1, \dots, n\}$, with ω being an $(n + 1)$ -root of 1, $\omega^{n+1} = 1$.

In the general case, the conjugacy class $[n_1, n_2, \dots, n_k]$ is simply the conjugacy class of the Coxeter element of the regular subalgebra (5.7) associated to the partition (5.6). By the Coxeter element of a semisimple Lie algebra we mean just the composition of the Coxeter elements of each simple ideal. Given a regular subalgebra of a simple Lie algebra g , the Cartan subalgebra of g can be constructed as the union of the Cartan subalgebras of each simple ideal of the subalgebra plus some additional elements to match the rank of g ; obviously, the Coxeter element of the regular subalgebra leaves these additional elements invariant. In the case of (5.7), the number of invariant directions is precisely $k - 1$, as follows

from the difference between the rank of the right-hand-side (n) and of the left-hand-side ($\sum_{j=1}^k (n_j - 1) = n + 1 - k$).

As summarized by (4.7) and (4.8), to construct the Heisenberg subalgebra of $\widehat{g}$ associated to a conjugacy class of the Weyl group of g , one needs a lift of a representative element of the class onto $\widehat{g}$; the lift of the Coxeter element is uniquely defined. For A_n , it can be expressed as

$$\widehat{w}_c = \exp \left(\frac{2\pi i}{2(n+1)} \text{ad } J_0^{(p)} \right), \quad (5.10)$$

where $J_0^{(p)}$ has been given in (5.3). The Heisenberg subalgebra of $\widehat{g}$ associated to the conjugacy class of the Coxeter element is called the principal Heisenberg subalgebra, and it can be defined as the centralizer in $\widehat{g}$ of the affine “cyclic element” of B. Kostant [28],

$$b^{(K)} = \sum_{j=1}^r \lambda_j 1 \otimes E_{\alpha_j} + \lambda_0 z \otimes E_{-\psi}, \quad (5.11)$$

where the λ_j 's are arbitrary non-vanishing coefficients, usually taken to be 1. $\psi = \sum_{j=1}^r k_j \alpha_j$ is the highest root of g , and $k_0 = 1$, $k_1, \dots$, and k_r are the Kac labels of $\widehat{g}$. Regarding A_n and using the fundamental representation of $sl(n+1, \mathbb{C})$, the affine cyclic element is the well known $(n+1) \times (n+1)$ matrix

$$\left(b^{(K)} \right)_{A_n} = \begin{pmatrix} 0 & 1 & & & \\ & 0 & 1 & & \\ & & \ddots & \ddots & \\ & & & 0 & 1 \\ z & & & & 0 \end{pmatrix}, \quad (5.12)$$

and the elements of the principal Heisenberg subalgebra are

$$b_{j+(n+1)m}^{(p)} = z^m \begin{pmatrix} 0 & \mathbb{I}_{n+1-j} \\ z \mathbb{I}_j & 0 \end{pmatrix}; \quad m \in \mathbb{Z}, \quad j = 1, \dots, n, \quad (5.13)$$

where $j + (n+1)m$ is the principal grade of this element, $\mathbb{I}_j$ is the $j \times j$ identity matrix, and $b_1^{(p)} = b^{(K)}$. Obviously, the associated gradation is also uniquely defined; it is the principal gradation $\mathbf{s}_p = (1, 1, \dots, 1)$ that is induced by $H_{\mathbf{s}_p} = J_0^{(p)}/2$.

We have already indicated that a conjugacy class $[w]$ of the Weyl group of A_n is the conjugacy class of the Coxeter element of some of its regular subalgebras, say $\widetilde{g}$. Therefore, a possible lift $\widehat{w}$ is obtained by composing the lifts of the Coxeter elements of the simple ideals of $\widetilde{g}$. For instance, for the conjugacy class $[w] = [n_1, n_2, \dots, n_k]$ the lift would be

$$\widehat{w} = \prod_{\substack{j=1 \\ n_j \neq 1}}^k \exp \left(\frac{2\pi i}{2n_j} \text{ad } \left(J_0^{(p)} \right)_{A_{n_j-1}} \right) \equiv \exp \left(\frac{2\pi i}{N_{\mathbf{s}_w}} \text{ad } H_{\mathbf{s}_w} \right), \quad (5.14)$$

where we have used the same notation as in (4.8). The order of $\widehat{w}$ is the following [40]: let $N'_{\mathbf{s}_w}$ be the least common multiple of $(n_1, n_2, \dots, n_k)$, then

$$N_{\mathbf{s}_w} = \begin{cases} N'_{\mathbf{s}_w}, & \text{if } N'_{\mathbf{s}_w} \left(\frac{1}{n_i} + \frac{1}{n_j} \right) \in 2\mathbb{Z} \text{ for any } i, j = 1, \dots, k; \\ 2N'_{\mathbf{s}_w}, & \text{otherwise.} \end{cases} \quad (5.15)$$

Moreover,

$$H_{\mathbf{s}_w} = \sum_{\substack{j=1 \\ n_j \neq 1}}^k \frac{N_{\mathbf{s}_w}}{2n_j} \left(J_0^{(\mathbf{p})} \right)_{A_{n_j-1}}. \quad (5.16)$$

The decomposition of $H_{\mathbf{s}_w}$ with respect to the basis (5.4) of the Cartan subalgebra would be

$$H_{\mathbf{s}_w} = \sum_{j=1}^{n+1} \gamma_j H_j, \quad \sum_{j=1}^{n+1} \gamma_j = 0; \quad (5.17)$$

the γ_j 's can be arranged such that $\gamma_{n+1} \leq \gamma_n \leq \dots \leq \gamma_1$, which corresponds to a change of the basis of simple roots similar to the one used to order the components of the defining vectors (5.2) of the embeddings of A_1 into A_n . Once ordered, they have the symmetry property $\gamma_j = -\gamma_{n+2-j}$, and it follows from (5.5) that $\gamma_1 = -\gamma_{n+1} = N_{\mathbf{s}_w}(n_k - 1)/2n_k$, where n_k is the maximum of $n_1, \dots, n_k$. $H_{\mathbf{s}_w}$ induces an integer gradation of $A_n^{(1)}$, $\mathbf{s}_w$, whose components are

$$\begin{aligned} (\mathbf{s}_w)_j &= \gamma_j - \gamma_{j+1} = (\mathbf{s}_w)_{n+1-j}, \quad j = 1, \dots, n \\ (\mathbf{s}_w)_0 &= N_{\mathbf{s}_w} - \sum_{j=1}^n (\mathbf{s}_w)_j = N_{\mathbf{s}_w} - 2\gamma_1 = \frac{N_{\mathbf{s}_w}}{n_k}. \end{aligned} \quad (5.18)$$

For example, the gradation of $A_3^{(1)}$ associated to the conjugacy class $[1, 1, 2]$ is $(2, 1, 0, 1)$, while the gradation of $A_6^{(1)}$ associated to $[1, 2, 4]$ is $(2, 1, 1, 1, 1, 1)$; it is useful to compare these gradations with the “characteristics” of the embeddings associated to the same partitions.

The Heisenberg subalgebra associated to the conjugacy class $[w] = [n_1, n_2, \dots, n_k]$ is obtained as follows. Consider the Cartan subalgebra defined by joining the Cartan subalgebras of each simple factor in (5.7) plus the required $k - 1$ additional generators. The corresponding Heisenberg subalgebra is just the affinization of this Cartan subalgebra according to the automorphism (5.14). Then, it will be generated by the linear combinations of the principal Heisenberg subalgebras associated to each of the simple factors in (5.7), embedded in A_n , plus the affinization of the $k - 1$ additional generators (see also [19, 40]).

As a practical example, we present all the inequivalent Heisenberg subalgebras of $A_2^{(1)}$ and $A_3^{(1)}$ in the appendix. By construction, the Heisenberg subalgebra has a basis whose generators are graded by the gradation $\mathbf{s}_w$ defined in (5.18), and their grades follow from those of the principal Heisenberg subalgebras of each simple ideal in (5.7). They are⁶

$$\underbrace{0, \dots, 0}_{k-1 \text{ times}}, \bigcup_{\substack{j=1 \\ n_j \neq 1}}^k \left\{ \frac{N_{\mathbf{s}_w}}{n_j}, \frac{2N_{\mathbf{s}_w}}{n_j}, \dots, \frac{(n_j-1)N_{\mathbf{s}_w}}{n_j} \right\} \bmod (N_{\mathbf{s}_w}), \quad (5.19)$$

which are obviously integers because of (5.15). Notice that, also by construction, the $H_{\mathbf{s}_w}$ of the lift (5.14) satisfies the condition (4.9).

It follows from (5.19) that the lowest positive grade of the generators of the Heisenberg subalgebra of $A_n^{(1)}$ associated to the conjugacy class $[w] = [n_1, n_2, \dots, n_k]$ is $N_{\mathbf{s}_w}/n_k$, which, taking into account (5.18), agrees with the result of lemma 6. Moreover, it also follows from (5.3), (5.8), and (5.11) that the generator J_+ of the embedding of A_1 associated to the partition $n+1 = n_1 + \dots + n_k$ can be expressed as

$$J_+^{(n_k, \dots, n_1)} = \sum_{\substack{j=1 \\ n_j \neq 1}}^k \left(\left(b_1^{(\mathbf{p})} \right)_{A_{n_j-1}} \right)_0, \quad (5.20)$$

where $(\)_0$ indicates the projection of the components whose homogeneous grade is zero.

The eqs.(5.16) and (5.20) specify some sort of canonical relation between the non-conjugated embeddings of A_1 into A_n and the non-equivalent Heisenberg subalgebras of $A_n^{(1)}$, a relation which is possible to generalize for the other classical simple Lie algebras [42]. Nevertheless, it only involves the particular elements of the Heisenberg subalgebra directly related to the affine cyclic element of Kostant. But, in general, one needs to relate the zero-grade component of any element of an arbitrary Heisenberg subalgebra to the A_1 subalgebras. In the case of $A_n^{(1)}$ it is possible to do it by using the explicit representation (5.13). Let us take $n \geq 2$ and $l = 1, \dots, n-1$, then $n = ml + r$ for some $0 \leq r < l$, where m is the integer part of the ratio n/l . Next, consider the nilpotent element $\left(b_l^{(\mathbf{p})} \right)_0 \in A_{n-1}$ as the J_+ of an A_1 subalgebra of A_{n-1} . After a tedious calculation, one can show that J_0

⁶ Notice that this essentially agrees with the results of ref.[29] except that the possibility of $N_{\mathbf{s}_w}$ being twice the least common multiple of $\{n_1, \dots, n_k\}$ has not been considered there. Then, the condition to have a grade 1 element in $\mathcal{H}[w]$ is that $N_{\mathbf{s}_w} = n_k$, the maximum of $n_1, \dots, n_k$. This requires not only that $n_k/n_j \in \mathbb{Z}^+$ for any $j = 1, \dots, k$, as already realized in [29], but also that either all the (n_k/n_j) 's are even or that all of them are odd.

is the matrix [10,43]

$$J_0 = \text{diag} \left(\underbrace{m, \dots, m}_{r \text{ times}}, \underbrace{m-1, \dots, m-1}_{(l-r) \text{ times}}, \dots, \underbrace{-m, \dots, -m}_{r \text{ times}} \right), \quad (5.21)$$

where the multiplicities r and $(l-r)$ occur alternately, ending always with r . Let us point out that, within this representation of A_{n-1} , the main diagonal of J_0 is precisely the defining vector of the embedding; therefore, using (5.5), one can prove the following result.

Lemma 8. *The nilpotent element, $(b_l^{(p)})_0 \in A_{n-1}$, where $n = ml + r \geq 2$ with $l \geq 1$, $0 \leq r < l$, and $m \in \mathbb{Z} > 0$, can be considered as the J_+ of the A_1 subalgebra of A_{n-1} associated to the partition*

$$n = \underbrace{(m+1) + \dots + (m+1)}_{r \text{ times}} + \underbrace{m + \dots + m}_{(l-r) \text{ times}}, \quad (5.22)$$

i.e., to the regular subalgebra

$$(A_m)^{\oplus r} \oplus (A_{m-1})^{\oplus (l-r)} \subset A_{n-1}. \quad \square \quad (5.23)$$

This result leads to the generalization of (5.20) in an obvious way, but we shall omit the explicit formula because it becomes too cumbersome. In any case, this shows that there are different forms of expressing the J_+ of an A_1 subalgebra of A_n as the zero graded component of some element of a Heisenberg subalgebra of $A_n^{(1)}$.

The following result of [43] will be used in the next section.

Lemma 9. *Let B_- be the maximal nilpotent negative subalgebra of A_{n-1} ; i.e., it corresponds to the lower triangular $n \times n$ matrices in the fundamental representation of $sl(n, \mathbb{C})$. Then*

$$\text{Ker} \left(\text{ad} \left(b_l^{(p)} \right)_0 \right) \cap B_- = \emptyset \quad (5.24)$$

if and only if either $l = 1$, or $l = 2$ and n is odd. $\square$

Finally, let us mention the following result of [19] that is relevant in relation to the distinction between hierarchies of type I and type II —recently, this result has been generalized for the other classical Lie algebras in [25].

Proposition 4. (Theorem 1.7 of [19]) *Graded regular elements exist only in those Heisenberg subalgebras of $A_n^{(1)}$ which belong to the special partitions*

$$n+1 = \underbrace{a + \cdots + a}_{p \text{ times}} \quad \text{and} \quad n+1 = \underbrace{a + \cdots + a}_{p \text{ times}} + 1,$$

where $a > 1$. The graded regular elements are of the form

$$\Lambda = z^m \left(\sum_{j=1}^p y_j \left(b_l^{(p)} \right)_{A_{a-1}} \right), \quad (5.25)$$

where

$$1 \leq l \leq a-1, \quad y_i \neq 0, \quad y_i^a \neq y_j^a, \quad i, j = 1, \dots, p, \quad i \neq j,$$

l is relatively prime to a , and $m \in \mathbb{Z}$. When

$$n+1 = \underbrace{a + \cdots + a}_{p \text{ times}} + 1$$

and a is even, the element Λ has grade $2(l+ma)$ with respect to the gradation (5.18), where $N_{\mathbf{s}_w} = 2a$. Otherwise, Λ is of grade $(l+ma)$ and $N_{\mathbf{s}_w} = a$. $\square$

6. $\mathcal{W}$ -algebras and integrable hierarchies of the KdV type.

We are now in position to investigate which are the $\mathcal{W}$ -algebras arising as the second Poisson bracket algebra of the integrable hierarchies of [21,22] in the sense of theorem 2. In the following, we restrict ourselves to the generalized KdV hierarchies based on $A_n^{(1)}$, for which $\mathbf{s}$ is the homogeneous gradation $\mathbf{s}_h = (1, 0, \dots, 0)$.

Theorem 2 indicates that, under (3.8), the second Poisson bracket algebra of the integrable hierarchy associated to the data $\{\widehat{g}, \mathcal{H}[w], \mathbf{s}_w, \mathbf{s}, \Lambda\}$ contains the $\mathcal{W}$ -algebra characterized by $J_+ = (\Lambda)_0$. Therefore, it will be possible to associate at least one integrable hierarchy to those $\mathcal{W}$ -algebras corresponding to the A_1 subalgebras of A_n such that J_+ can be expressed as the zero homogeneous grade component of an element of some Heisenberg subalgebra with definite $\mathbf{s}_w$ -grade. The results of the previous section allow us to express the J_+ of any A_1 subalgebra of A_n as the zero homogeneous grade component of, at least,

one element of some Heisenberg subalgebra. Accordingly, the condition will be that at least one of these elements have definite $\mathbf{s}_w$ -grade.

Let us analyse first the canonical connection expressed by (5.20). With respect to the gradation $\mathbf{s}_w$ induced by (5.16), the cyclic element $(b^{(K)})_{A_{n_j-1}}$ has grade $N_{\mathbf{s}_w}/n_j$. Consequently, the condition that the element of the Heisenberg subalgebra on the right-hand-side of (5.20) has definite $\mathbf{s}_w$ -grade requires that all the $n_j \neq 1$ are equal. This restricts the choice of J_+ to partitions of the type

$$n + 1 = \underbrace{a + \dots + a}_{p \text{ times}} + \underbrace{1 + \dots + 1}_{q \text{ times}} \equiv p(a) + q(1), \quad (6.1)$$

with $a > 1$. With this restriction,

$$H_{\mathbf{s}_w} = \frac{N_{\mathbf{s}_w}}{2a} J_0^{(a, \dots, a, 1, \dots, 1)}, \quad (6.2)$$

where

$$[w] = [\underbrace{a, \dots, a}_{p \text{ times}}, \underbrace{1, \dots, 1}_{q \text{ times}}]. \quad (6.3)$$

Notice that $H_{\mathbf{s}_w}$ is proportional to the J_0 of the relevant A_1 subalgebra, which ensures that condition (3.8) is indeed satisfied —see comments below theorem 2. Moreover, according to lemma 4, the $\mathbf{s}_w$ -grade of Λ, i , is either 1 or 2.

The gradations associated to the partitions (6.1) are the following.

Lemma 10. *The gradation $\mathbf{s}_w$ corresponding to the conjugacy class (6.3) is*

$$\mathbf{s}_w = \left(\underbrace{1, 0_{p-1}, \dots, 1, 0_{p-1}}_{\frac{a+1}{2} \text{ times}}, 0_q, \underbrace{1, 0_{p-1}, \dots, 1, 0_{p-1}}_{\frac{a-1}{2} \text{ times}} \right), \quad (6.4)$$

with $N_{\mathbf{s}_w} = a$, when a is an odd number,

$$\mathbf{s}_w = \left(\underbrace{2, 0_{p-1}, \dots, 2, 0_{p-1}}_{\frac{a}{2} \text{ times}}, 1, 0_{q-1}, 1, 0_{p-1}, \underbrace{2, 0_{p-1}, \dots, 2, 0_{p-1}}_{\frac{a}{2}-1 \text{ times}} \right), \quad (6.5)$$

with $N_{\mathbf{s}_w} = 2a$, when a is even and $q \neq 0$, and

$$\mathbf{s}_w = \left(\underbrace{1, 0_{p-1}, \dots, 1, 0_{p-1}}_{a \text{ times}} \right), \quad (6.6)$$

with $N_{\mathbf{s}_w} = a$, when a is even and $q = 0$. In these equations we have used the notation $0_j = \underbrace{0, \dots, 0}_{j \text{ times}}$.

Proof. When a is odd, it follows from (5.15) that $N_{\mathbf{s}_w} = a$. Besides, (5.16) leads to

$$\begin{aligned} H_{\mathbf{s}_w} &\equiv (\gamma_1, \dots, \gamma_{n+1}) \\ &= \left(\left(\frac{a-1}{2} \right)_p, \left(\frac{a-3}{2} \right)_p, \dots, 1_p, 0_{p+q}, -1_p, \dots, -\left(\frac{a-1}{2} \right)_p \right), \end{aligned} \quad (6.7)$$

where we have already ordered the components of $H_{\mathbf{s}_w}$ as indicated below (5.17). Using (5.18), all this leads to (6.4). In a similar way, when a is even, $N_{\mathbf{s}_w} = 2a$, and

$$H_{\mathbf{s}_w} \equiv (\gamma_1, \dots, \gamma_{n+1}) = ((a-1)_p, (a-3)_p, \dots, 1_p, 0_q, -1_p, \dots, -(a-1)_p), \quad (6.8)$$

which proves (6.5) and (6.6). $\square$

Let us indicate that the $\mathbf{s}_w$ -grade of the element of $\mathcal{H}[w]$ that appears on the right-hand-side of (5.20) for the partition (6.1) is $i = (\mathbf{s}_w)_0$, *i.e.*, either $i = 1$ or $i = 2$ according to lemma 10; this is the lowest positive $\mathbf{s}_w$ -grade of the elements of $\mathcal{H}[w]$.

So far, we have only analysed the canonical relation (5.20) between the A_1 subalgebras of A_n and the Heisenberg subalgebras of $A_n^{(1)}$. We can conclude that it is only possible to derive the $\mathcal{W}$ -algebras associated to the partitions of the form (6.1) from the second Poisson bracket algebras constructed in terms of the Heisenberg subalgebras associated to the same partitions. Nevertheless, as we have already said in the previous section, there are more general connections between A_1 subalgebras and Heisenberg subalgebras that follow from lemma 8. Let us consider the conjugacy class

$$[w] = [n_1, \dots, n_k, \underbrace{1, \dots, 1}_{q \text{ times}}] \quad (6.9)$$

of the Weyl group of A_n , with $n_j \neq 1$ for $j = 1, \dots, k$, and the nilpotent element

$$J_+ = \left(\sum_{j=1}^k y_j \left(b_{l_j}^{(p)} \right)_{A_{n_j-1}} \right)_0, \quad (6.10)$$

with $1 \leq l_j < n_j$ for $j = 1, \dots, k$. In this case, the condition that the element of $\mathcal{H}[w]$ defining the right-hand-side of (6.10) has definite $\mathbf{s}_w$ -grade is accomplished if

$$\frac{N_{\mathbf{s}_w}}{n_1} l_1 = \frac{N_{\mathbf{s}_w}}{n_2} l_2 = \dots = \frac{N_{\mathbf{s}_w}}{n_k} l_k \quad (6.11)$$

which implies that

$$n_j = m l_j + r_j, \quad j = 1, \dots, k, \quad (6.12)$$

with $m \in \mathbb{Z} \geq 1$, $0 \leq r_j < l_j$, and either $r_j = 0$ for all $j = 1, \dots, k$, or $r_j \neq 0$ and $r_i/r_j = l_i/l_j$ for any $i, j = 1, \dots, k$. Hence, it follows from lemma 8 that the J_+ expressed by (6.10) corresponds to the embedding of A_1 associated to the partition

$$n = \underbrace{(m+1) + \dots + (m+1)}_{\sum_{j=1}^k r_j \text{ times}} + \underbrace{m + \dots + m}_{\sum_{j=1}^k (l_j - r_j) \text{ times}} + \underbrace{1 + \dots + 1}_{q \text{ times}}. \quad (6.13)$$

Let us impose now the condition (3.8). Our starting point is the observation that, because of eq.(5.16), the restriction

$$H_{\mathbf{s}_w} \Big|_{A_{n_j-1}} = \frac{N_{\mathbf{s}_w}}{2n_j} \left(J_0^{(p)} \right)_{A_{n_j-1}}, \quad (6.14)$$

which implies that

$$\bigoplus_{\substack{j=1 \\ n_j \neq 1}}^k (B_-)_{A_{n_j-1}} \subsetneq P = \widehat{g}_0(\mathbf{s}) \cap \widehat{g}_{<0}(\mathbf{s}_w), \quad (6.15)$$

where $(B_-)_{A_{n_j-1}}$ is the maximal negative nilpotent subalgebra of A_{n_j-1} . Therefore, lemma 9 can be applied to each factor $\left(b_{l_j}^{(p)} \right)_{A_{n_j-1}}$ in eq.(6.10), which requires that⁷

$$\text{either } l_j = 1, \text{ or } l_j = 2 \text{ and } n_j \text{ is odd for all } j = 1, \dots, k. \quad (6.16)$$

In general, because the left-hand-side of eq.(6.15) does not equal P , (6.16) is just a necessary condition to ensure that (3.8) is satisfied. Nevertheless, when combined with the condition that the element of $\mathcal{H}[w]$ on the right-hand-side of (6.10) has definite $\mathbf{s}_w$ -grade, it implies that all the n_j 's have to be equal; then, according to (5.16), $H_{\mathbf{s}_w}$ is proportional to the J_0 of the A_1 subalgebra corresponding to J_+ in (6.10), and, consequently the condition (3.8) is indeed satisfied in all these cases, as we have discussed after theorem 2.

In conclusion, eqs.(6.12), (6.13), and (6.16) specify the only $\mathcal{W}$ -algebras that can be derived as the second Poisson bracket algebra of the integrable hierarchies of [21,22] based on a constant element $\Lambda \in \mathcal{H}[w]$ that satisfies the condition (3.8). This, together with theorem 2, is the main result of this paper and we summarize it as follows.

⁷ We thank Laszlo Fehér for pointing out this fact to us [44].

Theorem 3. (i) Graded elements that satisfy condition (3.8) exist only in those Heisenberg subalgebras of $A_n^{(1)}$ associated to the special conjugacy classes

$$[w] = \underbrace{[m, \dots, m]}_{k \text{ times}}, \underbrace{[1, \dots, 1]}_{q \text{ times}}. \quad (6.17)$$

These elements are of the form

$$\Lambda^{(1)} = \sum_{j=1}^k y_j \left(b_1^{(p)} \right)_{A_{m-1}}, \quad (6.18)$$

for any value of m , and of the form

$$\Lambda^{(2)} = \sum_{j=1}^k y_j \left(b_2^{(p)} \right)_{A_{m-1}}, \quad (6.19)$$

when m is odd. If either m is odd or m is even and $q = 0$, the $\mathbf{s}_w$ -grade of $\Lambda^{(1)}$ equals 1; otherwise, $\Lambda^{(1)} \in \widehat{g}_2(\mathbf{s}_w)$. In any case, the $\mathbf{s}_w$ -grade of $\Lambda^{(1)}$ is the lowest positive $\mathbf{s}_w$ -grade of the elements of $\mathcal{H}[w]$. The $\mathbf{s}_w$ -grade of $\Lambda^{(2)}$ equals 2.

(ii) Consider now the integrable hierarchies of the KdV type associated to $A_n^{(1)}$, $\mathcal{H}[w]$, $\mathbf{s}_w$, $\mathbf{s}_h = (1, 0, \dots, 0)$, and to one of those two elements of $\mathcal{H}[w]$. The algebras defined by the second Poisson bracket of these integrable hierarchies, restricted to the gauge invariant functionals of $q_0(x)$, are the $\mathcal{W}$ algebras associated to the embeddings of A_1 into A_n labelled by the partitions

$$n = \underbrace{m + \dots + m}_{k \text{ times}} + \underbrace{1 + \dots + 1}_{q \text{ times}} \equiv k(m) + q(1), \quad (6.20)$$

for $\Lambda^{(1)}$, or

$$\begin{aligned} n &= \underbrace{(a+1) + \dots + (a+1)}_{k \text{ times}} + \underbrace{a + \dots + a}_{k \text{ times}} + \underbrace{1 + \dots + 1}_{q \text{ times}} \\ &\equiv k(a+1) + k(a) + q(1), \end{aligned} \quad (6.21)$$

for $\Lambda^{(2)}$ when m is odd, $m = 2a + 1$. $\square$

Remark. Using proposition 4, one can restrict theorem 3 to the integrable hierarchies of type I. Then, q has to be either 0 or 1, and $y_i^m \neq y_j^m$ for any $i \neq j = 1, \dots, k$.

Notice that the connection expressed by theorem 3 shows that the $\mathcal{W}$ -algebras specified by the partitions $n = k(2) + q(1)$ are shared as second Poisson bracket algebras by the

integrable hierarchies associated to the conjugacy classes

$$[w] = \underbrace{[2, \dots, 2]}_{k \text{ times}}, \underbrace{[1, \dots, 1]}_{q \text{ times}}, \quad \text{with } \Lambda^{(1)} \in \mathcal{H}[w], \quad (6.22)$$

and

$$[w] = \underbrace{[3, \dots, 3]}_{k \text{ times}}, \underbrace{[1, \dots, 1]}_{q-k \text{ times}}, \quad \text{with } \Lambda^{(2)} \in \mathcal{H}[w]. \quad (6.23)$$

Nevertheless, in general, the $\mathcal{W}$ -algebras associated to the partitions (6.20) or (6.21) can be obtained only from integrable hierarchies where Λ has to be of the form $\Lambda^{(1)}$ or $\Lambda^{(2)}$, respectively.

Finally, let us point out that the class of $\mathcal{W}$ -algebras covered by theorems 2 and 3 is very restricted even if type II hierarchies are considered; a fact that has been already anticipated in [19]. In fact, notice that the structure of the second Poisson bracket algebra of the integrable hierarchies is actually independent of the fact that they are of type I or of type II as far as (2.6) is satisfied. The first $\mathcal{W}$ -algebra that cannot be recovered as the second Poisson bracket algebra of any of the integrable hierarchies of [21,22], with Λ restricted by the condition (3.8), corresponds to the partition $6 = 4 + 2$, *i.e.*, to the regular subalgebra $A_3 \oplus A_1 \subset A_5$.

7. Examples.

7.1 The First Fractional A_2 KdV-Hierarchy and $\mathcal{W}_3^{(2)}$ again.

This paper has been largely motivated by the analysis of this case in [22]; here, we discuss it again to illustrate some points of the previous sections. Let us consider the KdV hierarchy corresponding to the principal Heisenberg subalgebra of $A_2^{(1)}$ and to the element Λ with principal grade $i = 2$,

$$\Lambda = \begin{pmatrix} 0 & 0 & 1 \\ z & 0 & 0 \\ 0 & z & 0 \end{pmatrix}. \quad (7.1)$$

Since this Λ is not the element of the principal Heisenberg subalgebra with minimal positive grade, this integrable hierarchy can be considered just as the result of changing x by the

first time of the generalized A_2 Drinfel'd-Sokolov hierarchy [14,15]. Before gauge fixing, the potential can be written as

$$q(x) = \begin{pmatrix} y_1(x) & c(x) & 0 \\ e(x) & y_2(x) & d(x) \\ a(x) + z b(x) & f(x) & -(y_1(x) + y_2(x)) \end{pmatrix} \quad (7.2)$$

which, under a gauge transformation, becomes

$$q(x) \mapsto \Phi(x) \partial_x \Phi^{-1}(x) + \Phi(x) (\Lambda + q(x)) \Phi^{-1}(x) - \Lambda ,$$

where

$$\Phi(x) = \exp S(x) = \begin{pmatrix} 1 & 0 & 0 \\ A(x) & 1 & 0 \\ B(x) & C(x) & 1 \end{pmatrix} . \quad (7.3)$$

First of all, let us identify the A_1 subalgebra of A_2 corresponding to this case, whose generators are

$$J_+ = (\Lambda)_0 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} , \quad J_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} , \quad \text{and} \quad J_- = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} . \quad (7.4)$$

According to Section 5, the form of J_0 indicates that it is the A_1 subalgebra corresponding to the partition $3 = 2 + 1$. Moreover, the principal gradation is induced precisely by $H_{\mathbf{s}_w} = J_0$ which obviously fulfills the condition (3.8). Now, instead of fixing the gauge as in [21,22], we choose the gauge slice according to theorem 1. Then,

$$q^{(\text{can})}(x) = 3\phi(x) \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ z & 0 & 0 \end{pmatrix} + \begin{pmatrix} U(x)/2 & 0 & 0 \\ G^+(x) & -U(x) & 0 \\ T(x) & G^-(x) & U(x)/2 \end{pmatrix} , \quad (7.5)$$

which leads to the functional dependence on $q_0(x)$ of the components of $\widehat{S}(x)$ (or equivalently $\widehat{\Phi}(x)$)

$$\widehat{A} = -d , \quad \widehat{C} = c , \quad \widehat{B} = \frac{1}{2} (-cd + 2y_1 + y_2) , \quad (7.6)$$

and of the components of $\mathcal{J}(x)$

$$\begin{aligned} \phi &= \frac{1}{3} (b + c + d) \\ U &= cd - y_2 \\ G^+ &= -cd^2 + e - d(y_1 - y_2) + d' \\ G^- &= -c^2d + f + c(y_1 + 2y_2) - c' . \end{aligned} \quad (7.7)$$

The energy-momentum tensor (4.3) is

$$\begin{aligned}\mathcal{T}(x) &= T + \frac{3}{4}U^2 \\ &= a + ce + d f + y_1^2 + y_1 y_2 + y_2^2 \\ &\quad + \frac{1}{2}(cd' - dc') - y_1' - \frac{1}{2}y_2' .\end{aligned}\tag{7.8}$$

One can check that ϕ , U , $G^\pm$, and $\mathcal{T}$ are exactly equal to the combinations ϕ , $\tilde{U}$, $\tilde{G}^\pm$, and $\tilde{T}$ of [22], respectively. We already knew that, in this case, the restriction of the second Poisson bracket to the components of $q_0^{(\text{can})}(x)$ is a $\mathcal{W}$ -algebra; in fact, it is the $\mathcal{W}$ -algebra associated to the A_1 subalgebra of A_2 labelled by the partition $3 = 2 + 1$. This realization of this $\mathcal{W}$ -algebra is an example of the connection expressed by theorem 3, with $[w] = [3]$ and $\Lambda = \Lambda^{(2)}$.

But it also follows from theorem 3 that this $\mathcal{W}$ -algebra can also be realized as the second Poisson bracket of a different integrable hierarchy with $[w] = [2, 1]$ and $\Lambda = \Lambda^{(1)}$. This is the canonical connection expressed by (5.20), (6.1), and (6.3), which leads precisely to the realization discussed in [22] (see also [17]), with $J_+ = (\Lambda')_0$ and

$$\Lambda' = \Lambda^{(1)} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ z & 0 & 0 \end{pmatrix} \in \mathcal{H}[2, 1] .\tag{7.9}$$

Finally, let us recall that Λ and Λ' are regular elements of the corresponding Heisenberg subalgebras of $A_2^{(1)}$ and, hence, these integrable hierarchies are of type I.

7.2 Other Fractional A_{N-1} KdV-Hierarchies and $\mathcal{W}$ -algebras.

The example analysed in the previous section is just the simplest case of the so called fractional KdV-hierarchies [16]. Let us now consider the KdV general hierarchy corresponding to the principal Heisenberg subalgebra of $A_{N-1}^{(1)}$, $[w] = [N]$, and to the element Λ with principal grade $i = 2$, *i.e.*,

$$\Lambda = b_2^{(\text{p})} = \begin{pmatrix} 0 & \mathbb{I}_{(N-2) \times (N-2)} \\ z \mathbb{I}_{2 \times 2} & 0 \end{pmatrix} .\tag{7.10}$$

According to lemma 8, the embedding of A_1 into A_{N-1} corresponding to $J_+ = \left(b_2^{(\text{p})}\right)_0$ is labelled by the partition

$$N = \left(\frac{N+1}{2}\right) + \left(\frac{N-1}{2}\right)\tag{7.11}$$

when $N \geq 3$ is odd, and by the partition

$$N = \left(\frac{N}{2}\right) + \left(\frac{N}{2}\right) \quad (7.12)$$

when $N \geq 4$ is even. Nevertheless, before claiming any relation between the second Poisson bracket of this hierarchy and $\mathcal{W}$ -algebras, one has to check if the condition (3.8) is satisfied.

Let us start with the case when N is odd. Then, it follows from lemma 9 that indeed the condition (3.8) holds in this case. Then, the second Poisson bracket algebra of this KdV hierarchy contains the $\mathcal{W}$ -algebra associated to the partition (7.11), which is nothing else than the fractional $\mathcal{W}_N^{(2)}$ algebra of [45,43]. This case is the straightforward generalization of the first example that we have analysed in this section, which is recovered when $N = 3$.

Let us now study the case when N is even. Then, it is easy to check that

$$\theta = \sum_{j=1}^{N/2} E_{2j,2j-1} \in P = g_0(\mathbf{s}) \cap g_{<0}(\mathbf{s}_w) \quad (7.13)$$

belongs to $\text{Ker}(\text{ad } J_+)$ as well; hence, the condition (3.8) is not satisfied in agreement with lemma 9. Even more, one can check that $[\Lambda, \theta] = 0$; therefore, neither (3.8) nor (2.6) are satisfied, and one cannot construct an integrable hierarchy in the sense of eq.(2.5) in this case.

Let us now go on to consider the KdV hierarchy corresponding to the element Λ with principal grade $i = 3$, *i.e.*,

$$\Lambda = b_3^{(p)} = \begin{pmatrix} 0 & \mathbb{I}_{(N-3) \times (N-3)} \\ z \mathbb{I}_{3 \times 3} & 0 \end{pmatrix}. \quad (7.14)$$

The A_1 subalgebra of A_N with $J_+ = (b_2^{(p)})_0$ is labelled by the partition

$$N = \left(\frac{N+2}{3}\right) + \left(\frac{N-1}{3}\right) + \left(\frac{N-1}{3}\right) \quad (7.15)$$

when $N \in 3\mathbb{Z} + 1 \geq 4$, by the partition

$$N = \left(\frac{N+1}{3}\right) + \left(\frac{N+1}{3}\right) + \left(\frac{N-2}{3}\right), \quad (7.16)$$

when $N \in 3\mathbb{Z} + 2 \geq 5$, and by the partition

$$N = \left(\frac{N}{3}\right) + \left(\frac{N}{3}\right) + \left(\frac{N}{3}\right), \quad (7.17)$$

when $N \in 3\mathbb{Z} \geq 6$.

It is easy to check that

$$\theta_1 = \sum_{j=1}^{(N-1)/3} E_{3j,3j-1} \in P \quad (7.18)$$

belongs to $\text{Ker}(\text{ad } J_+)$ when $N \in 3\mathbb{Z} + 1 \geq 4$, that

$$\theta_2 = \sum_{j=1}^{(N+1)/3} E_{3j-1,3j-2} \in P \quad (7.19)$$

belongs to $\text{Ker}(\text{ad } J_+)$ when $N \in 3\mathbb{Z} + 2 \geq 5$, and that

$$\theta_3 = \sum_{j=1}^{N/3} (\alpha E_{3j,3j-1} + \beta E_{3j-1,3j-2} + \gamma E_{3j,3j-2}) \in P \quad (7.20)$$

belongs to $\text{Ker}(\text{ad } J_+)$ when $N \in 3\mathbb{Z} \geq 6$ for arbitrary constants α , β , and γ . Thus, we conclude that the condition (3.8) is not satisfied in this case, as anticipated by lemma 9. Nevertheless, one can notice that $[\Lambda, \theta_3] = 0$, but $[\Lambda, \theta_1]$ and $[\Lambda, \theta_2]$ do not vanish. Hence, (2.6) holds except when $N \in 3\mathbb{Z} \geq 6$ and one can indeed construct integrable hierarchies in all the other cases. It is expected that their second Poisson bracket algebras will contain $\mathcal{W}$ -algebras as well, but they are not described by the results of this paper (see [15,18]).

Finally, let us point out that all these fractional A_{N-1} hierarchies can be described as the result of changing x for some of the times t_b of the generalized A_{N-1} Drinfel'd-Sokolov hierarchy, in the sense of [14,15], only when $b \in \mathcal{H}[w]$ is regular and the resulting hierarchy is of type I.

7.3 A Fractional KdV-Hierarchy associated to $\mathcal{H}[3,3] \subset A_5^{(1)}$.

The last example is an integrable hierarchy such that $q_{>0}^{(\text{can})}(x)$ has dynamical components, *i.e.*, it has some components that are not centres of the second Poisson bracket. Within the $A_n^{(1)}$ algebras, the simplest case is associated to the Heisenberg subalgebra of $A_5^{(1)}$ corresponding to the conjugacy class of the Weyl group $[w] = [3,3]$.

According to (5.15), $N_{\mathbf{s}_w} = 3$, and, using (5.16), one gets (see the appendix),

$$H_{\mathbf{s}_w} = \text{diag} \left(\frac{3}{2 \cdot 3}(2, 0, -2) \cup \frac{3}{2 \cdot 3}(2, 0, -2) \right) \mapsto \text{diag} (1, 1, 0, 0, -1, -1), \quad (7.21)$$

and the gradation associated to $\mathcal{H}[3, 3]$ is

$$\mathbf{s}_w = (1, 0, 1, 0, 1, 0) . \quad (7.22)$$

Using the basis of simple roots associated to $\mathbf{s}_w$, the generators of the Heisenberg subalgebra are

$$b_{3m} = z^m \text{diag} (1, -1, 1, -1, 1, -1) \quad (7.23)$$

$$b_{1+3m}^{(1)} = z^m \begin{pmatrix} & & 0 & & & \\ & & 1 & & & \\ & & & 0 & & \\ & & & & 1 & \\ & & & & & 1 \\ 0 & & & & & z \end{pmatrix} \quad b_{1+3m}^{(2)} = z^m \begin{pmatrix} & & & & 1 & \\ & & & & 0 & \\ & & & & & 1 \\ & & & & & & 0 \\ z & & & & & & \\ & 0 & & & & & \end{pmatrix} \quad (7.24)$$

$$b_{2+3m}^{(1)} = z^m \begin{pmatrix} & & & & 0 & \\ & & & & & 1 \\ 0 & & & & & \\ & z & & & & \\ & & 0 & & & \\ & & & z & & \end{pmatrix} \quad b_{2+3m}^{(2)} = z^m \begin{pmatrix} & & & & 1 & \\ & & & & & 0 \\ z & & & & & \\ & 0 & & & & \\ & & z & & & \\ & & & 0 & & \end{pmatrix}, \quad (7.25)$$

where $m \in \mathbb{Z}$ and the subscripts are the $\mathbf{s}_w$ -grades.

Let us consider the type I hierarchy associated to $A_5^{(1)}$, $\mathcal{H}[3, 3]$, $\mathbf{s}_{[3,3]}$, $\mathbf{s}_h = (1, 0, \dots, 0)$, and the regular element

$$\Lambda = \begin{pmatrix} & & & & 1 & \\ & & & & & 2 \\ z & & & & & \\ & 2z & & & & \\ & & z & & & \\ & & & 2z & & \end{pmatrix} = b_2^{(2)} + 2b_2^{(1)} \in \mathcal{H}[3, 3], \quad (7.26)$$

whose grade is $i = 2$. In block form, the potential is

$$q(x) = \begin{pmatrix} \mathbb{A}(x) & \mathbb{B}(x) & 0 \\ \mathbb{C}(x) & \mathbb{D}(x) & \mathbb{E}(x) \\ \mathbb{F}(x) & \mathbb{G}(x) & \mathbb{H}(x) \end{pmatrix} + z \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \mathbb{J}(x) & 0 & 0 \end{pmatrix} \in \mathcal{Q}, \quad (7.27)$$

where all the entries are 2×2 matrices, and the gauge transformations are generated by elements of the form

$$S(x) = \begin{pmatrix} 0 & 0 & 0 \\ \mathbb{T}(x) & 0 & 0 \\ \mathbb{U}(x) & \mathbb{V}(x) & 0 \end{pmatrix} \in P; \quad (7.28)$$

$q(x)$ has 35 independent components, while $S(x)$ has 12. According to theorem 1, the gauge slice can be chosen such that

$$q^{(\text{can})}(x) = \begin{pmatrix} a_1 & a_2 & 0 & 0 & 0 & 0 \\ 2a_3 & -a_4 & 0 & 0 & 0 & 0 \\ b_1 & b_2 & a_5 - 2a_1 & a_6 & 0 & 0 \\ b_3 & b_4 & a_7 & 2a_4 - a_5 & 0 & 0 \\ c_1 & c_2 & b_5 & b_6 & a_1 & 2a_2 \\ c_3 & c_4 & b_7 & b_8 & a_3 & -a_4 \end{pmatrix} + z \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ \phi & B & 0 & 0 & 0 & 0 \\ A & \chi & 0 & 0 & 0 & 0 \end{pmatrix}, \quad (7.29)$$

i.e., it has $23(= 35 - 12)$ independent components.

It follows from theorem 3, with $[w] = [3, 3]$ and $\Lambda = \Lambda^{(2)}$, *i.e.*, with $m = 3$, $k = 2$, and $q = 0$, that $\{a_1, \dots, a_7, b_1, \dots, b_8, c_1, \dots, c_4\}$ generate the $\mathcal{W}$ -algebra associated to the embedding of A_1 into A_5 labelled by the partition $6 = (2) + (2) + (1) + (1)$. The conformal structure is defined by

$$\mathcal{T}(x) = c_1 + 2c_4 + 3(a_1^2 + a_4^2) + a_5^2 - 2(a_1 + a_4)a_5 + 4a_2a_3 + a_6a_7, \quad (7.30)$$

and the generators $\{a_1, \dots, a_7\}$, $\{b_1, \dots, b_8\}$, and $\{c_1 - 2c_4, c_2, c_3\}$ have conformal spin 1, $3/2$, and 2, respectively. Moreover, in analogy with the case of $\mathcal{W}_3^{(2)}$, notice that this $\mathcal{W}$ -algebra could also be obtained from another integrable hierarchy associated to the Heisenberg subalgebra $\mathcal{H}[2, 2, 1, 1]$ and to $\Lambda = \Lambda^{(1)}$.

In contrast, the second Poisson brackets of the components of $q_1^{(\text{can})}(x)$ are not given by theorems 2 and 3. It is easy to check that $\phi(x)$ and $\chi(x)$ are the two centres associated by proposition 1 to $b_1^{(1)}$ and $b_1^{(2)}$, and, hence, they are not dynamical quantities. The brackets involving the other two components, $A(x)$ and $B(x)$, can be calculated with the help of the computer. The result is that the only non-zero second Poisson bracket is

$$\{A(x), B(y)\}_2 = \frac{7}{2} \delta(x - y), \quad (7.31)$$

and we conclude that $A(x)$ and $B(x)$ generate a “ b - c ” algebra that is decoupled from the $\mathcal{W}$ -algebra generated by the components of $q_0^{(\text{can})}(x)$. Therefore, in this case, it is

actually possible to write down an energy-momentum tensor that generates the conformal transformation of all the components of $q^{(\text{can})}(x)$

$$\widehat{\mathcal{T}}(x) = \mathcal{T}(x) + \frac{1}{7} \left(A(x)B'(x) - A'(x)B(x) \right), \quad (7.32)$$

which gives conformal spin $\frac{1}{2}$ to $A(x)$ and $B(x)$.

8. Conclusions and discussion.

We have presented a systematic study of the second Hamiltonian structures of the integrable hierarchies defined in [21,22] in terms of (untwisted) affine Kac-Moody algebras. They are based on a graded element Λ of some Heisenberg subalgebra, and we have restricted ourselves to the case when $\text{Ker}(\text{ad}(\Lambda)_0) \cap \widehat{\mathfrak{g}}_{<0}(\mathbf{s}_w) = \emptyset$. When $(\Lambda)_0 \neq 0$, the second Poisson bracket algebra contains one of the (classical) $\mathcal{W}$ -algebras defined through the method of Drinfel'd-Sokolov Hamiltonian reduction in [8,9,10]. We have also generalized some of the results of [21,22] for the type II hierarchies, but, even considering both type I and type II hierarchies, the class of $\mathcal{W}$ -algebras that can be obtained in this way is very limited. The restriction comes both from the condition that Λ is a *graded* element and from the condition that it satisfies $\text{Ker}(\text{ad}(\Lambda)_0) \cap \widehat{\mathfrak{g}}_{<0}(\mathbf{s}_w) = \emptyset$. It implies that, in the case of $A_n^{(1)}$ algebras, only the $\mathcal{W}$ -algebras based on the embeddings of A_1 into A_n labelled by the partitions of the form

$$n+1 = k(m) + q(1) \quad \text{and} \quad n+1 = k(m+1) + k(m) + q(1)$$

can be defined as the second Poisson bracket algebras of some of these integrable hierarchies. A relevant feature is that some of these $\mathcal{W}$ -algebras can be recovered from more than one integrable hierarchy. Notice that there is a well defined “Miura map” within each integrable hierarchy, and that the Miura map usually leads to the realization of the $\mathcal{W}$ -algebra in terms of free fields [5]. Therefore, there might be more than one possible realizations suggested by this connection.

Let us point out that our results cover most of the $\mathcal{W}$ -algebras that have already been related to integrable hierarchies of the KdV type in the literature. An example of the other ones are the $\mathcal{W}_4^{(2)}$ and $\mathcal{W}_4^{(3)}$ algebras of [18] and [15], respectively. The problem is that, in these cases, either the condition (2.6) is not satisfied, or it is not possible to

restrict the second Poisson bracket algebra to functions that only depend on $q_0(x)$. We think that a further generalization of our results to cover these cases can be achieved by comparing directly the second Poisson bracket with the Dirac bracket of the Hamiltonian reduction; this work is already in progress. More recently, a series of $\mathcal{W}$ -algebras associated to a new generalization of the Gel'fand and Dickey's pseudo-differential approach has been constructed in ref.[20]. The relation of these $\mathcal{W}$ -algebras to the Drinfel'd-Sokolov Hamiltonian reduction and, of course, to our approach is unclear yet. It would be very interesting to understand better this connection not only from the point of view of the classification of $\mathcal{W}$ -algebras, but also in order to have an alternative definition of the integrable hierarchies of [21,22]. In fact, it has already been shown in [19,25] that some of them can be defined using a matrix version of the Gel'fand–Dickey's pseudo-differential approach.

Finally, let us comment briefly about the possible applications of our results to the study of two-dimensional integrable field theories. It has already been said in the introduction that the $\mathcal{W}$ -algebras are naturally related to the (conformal) Toda field theories. Therefore, our results should help to a better understanding of the relationship between those conformal Toda field theories and the integrable hierarchies of the KdV type. Some steps in this direction have already been taken in ref.[30]. Even more, it is possible to associate a generalization of the Toda equation to the equations of the mKdV type [4,46]. In some cases, this generalized Toda equation can be understood as the equation of motion of a massive affine Toda field theory, and all this could clarify the connection between conformal and massive Toda models. In relation to this, we expect that our detailed discussion of the A_1 subalgebras of a simple Lie algebra and the Heisenberg subalgebras of its untwisted affinization will be particularly useful [29].

Appendix.

We present the non-equivalent Heisenberg subalgebras of $A_2^{(1)}$ and $A_3^{(1)}$ (see also [19] and [40]). We shall realize A_n in terms of the fundamental representation of $sl(n+1, \mathbb{C})$; then, the basis (5.4) is just

$$H_i = E_{i,i} , \quad E_{(e_j - e_k)} = E_{k,j} , \quad (\text{A.1})$$

where $E_{j,k}$ is the $(n+1) \times (n+1)$ matrix whose only non-vanishing entry is a 1 at the

position (j, k) . Here, we do not mention neither the principal Heisenberg subalgebra, associated to $[n+1]$, nor the homogeneous Heisenberg subalgebra, associated to $[1, \dots, 1]$.

A₂ :

- $[w] = [2, 1]$ ($A_1 \subset A_2$): $N_{\mathbf{s}_w} = 4$, and

$$H_{\mathbf{s}_w} = \text{diag} \left(\frac{4}{2 \cdot 2} (1, -1) \cup (0) \right) \mapsto \text{diag} (1, 0, -1), \quad (\text{A.2})$$

where the arrow indicates that we have ordered the components of $H_{\mathbf{s}_w}$; in this case, $\gamma_1 \mapsto \gamma_1$, $\gamma_2 \mapsto \gamma_3$, and $\gamma_3 \mapsto \gamma_2$. Then,

$$\mathbf{s}_w = (2, 1, 1), \quad \text{and} \quad g_0(\mathbf{s}_w) = u(1)^2. \quad (\text{A.3})$$

The generators of the Heisenberg subalgebra are

$$\begin{aligned} b_{4m} &= z^m \begin{pmatrix} 1 & & \\ & 1 & \\ & & -2 \end{pmatrix} \mapsto z^m \begin{pmatrix} 1 & & \\ & -2 & \\ & & 1 \end{pmatrix}, \\ b_{2+4m} &= z^m \begin{pmatrix} 0 & 1 & \\ z & 0 & \\ & & 0 \end{pmatrix} \mapsto z^m \begin{pmatrix} & & 1 \\ & 0 & \\ z & & \end{pmatrix}, \quad m \in \mathbb{Z}, \end{aligned} \quad (\text{A.4})$$

where the arrow indicates the same permutation of components as in (A.2), and the subscripts are the grades with respect to $\mathbf{s}_w$. As we have explained in the main text, this permutation corresponds to a change of the basis of the simple roots through conjugation with the Weyl group, to ensure that $H_{\mathbf{s}_w}$ indeed induces a $\mathbb{Z}$ -gradation. Notice that the elements of $\mathcal{H}[w]$ have block-diagonal form before performing this permutation.

To illustrate the non-uniqueness of the gradation $\mathbf{s}_w$, let us point out that the generators b_{4m} and b_{2+4m} have definite grade with respect to any gradation of the form $\mathbf{s}_w' = m(s_1 + s_2, s_1, s_2)$ their grades being $2m(s_1 + s_2)$ and $(1 + 2m)(s_1 + s_2)$, respectively.

A₃ :

- $[w] = [3, 1]$ ($A_2 \subset A_3$): $N_{\mathbf{s}_w} = 3$, and

$$H_{\mathbf{s}_w} = \text{diag} \left(\frac{3}{2 \cdot 3} (2, 0, -2) \cup (0) \right) \mapsto \text{diag} (1, 0, 0, -1); \quad (\text{A.5})$$

then,

$$\mathbf{s}_w = (1, 1, 0, 1), \quad \text{and} \quad g_0(\mathbf{s}_w) = A_1 \oplus u(1)^2. \quad (\text{A.6})$$

The generators of the Heisenberg subalgebra are

$$\begin{aligned} b_{3m} &= z^m \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -3 \end{pmatrix} \mapsto z^m \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & -3 & \\ & & & 1 \end{pmatrix}, \\ b_{1+3m} &= z^m \begin{pmatrix} 0 & 1 & 0 & \\ 0 & 0 & 1 & \\ z & 0 & 0 & \\ & & & 0 \end{pmatrix} \mapsto z^m \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ z & 0 & 0 & 0 \end{pmatrix}, \\ b_{2+3m} &= z^m \begin{pmatrix} 0 & 0 & 1 & \\ z & 0 & 0 & \\ 0 & z & 0 & \\ & & & 0 \end{pmatrix} \mapsto z^m \begin{pmatrix} 0 & 0 & 0 & 1 \\ z & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & z & 0 & 0 \end{pmatrix}. \end{aligned} \quad (\text{A.7})$$

- $[w] = [2, 2]$ ($A_1 \oplus A_1 \subset A_3$): $N_{\mathbf{s}_w} = 2$,

$$H_{\mathbf{s}_w} = \text{diag} \left(\frac{2}{2 \cdot 2}(1, -1) \cup \frac{2}{2 \cdot 2}(1, -1) \right) \mapsto \text{diag} \left(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2} \right), \quad (\text{A.8})$$

$$\mathbf{s}_w = (1, 0, 1, 0), \quad g_0(\mathbf{s}_w) = A_1 \oplus A_1 \oplus u(1), \quad (\text{A.9})$$

and

$$\begin{aligned} b_{2m} &= z^m \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & -1 & \\ & & & -1 \end{pmatrix} \mapsto z^m \begin{pmatrix} 1 & & & \\ & -1 & & \\ & & 1 & \\ & & & -1 \end{pmatrix}, \\ b_{1+2m}^{(1)} &= z^m \begin{pmatrix} 0 & 1 & & \\ z & 0 & & \\ & & 0 & \\ & & & 0 \end{pmatrix} \mapsto z^m \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ z & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ b_{1+2m}^{(2)} &= z^m \begin{pmatrix} 0 & & & \\ & 0 & & \\ & & 0 & 1 \\ & & z & 0 \end{pmatrix} \mapsto z^m \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & z & 0 & 0 \end{pmatrix}, \end{aligned} \quad (\text{A.10})$$

where the superscripts distinguish the different generators of $\mathcal{H}[w]$ having the same $\mathbf{s}_w$ -grade.

- $[w] = [2, 1, 1]$ ($A_1 \subset A_3$): $N_{\mathbf{s}_w} = 4$,

$$H_{\mathbf{s}_w} = \text{diag} \left(\frac{4}{2 \cdot 2} (1, -1) \cup (0) \cup (0) \right) \mapsto \text{diag} (1, 0, 0, -1), \quad (\text{A.11})$$

$$\mathbf{s}_w = (2, 1, 0, 1), \quad g_0(\mathbf{s}_w) = A_1 \oplus u(1)^2, \quad (\text{A.12})$$

and

$$\begin{aligned} b_{4m}^{(1)} &= z^m \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & -2 & \\ & & & 0 \end{pmatrix} \mapsto z^m \begin{pmatrix} 1 & & & \\ & 0 & & \\ & & -2 & \\ & & & 1 \end{pmatrix}, \\ b_{4m}^{(2)} &= z^m \begin{pmatrix} 0 & & & \\ & 0 & & \\ & & -1 & \\ & & & 1 \end{pmatrix} \mapsto z^m \begin{pmatrix} 0 & & & \\ & 1 & & \\ & & -1 & \\ & & & 0 \end{pmatrix}, \\ b_{2+4m} &= z^m \begin{pmatrix} 0 & 1 & & \\ z & 0 & & \\ & & 0 & \\ & & & 0 \end{pmatrix} \mapsto z^m \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ z & 0 & 0 & 0 \end{pmatrix}. \end{aligned} \quad (\text{A.13})$$

For A_4 and A_5 we just show the gradations $\mathbf{s}_w$ and the grades of the elements of the corresponding Heisenberg subalgebras. First, for the conjugacy classes of the Weyl group of A_4 ,

Conjugacy class $[w]$	Gradation $\mathbf{s}_w$	Grades of $\mathcal{H}[w]$
$[4, 1]$	$(2, 2, 1, 1, 2)$	$(0, 2, 4, 6) \bmod 8$
$[3, 2]$	$(4, 1, 3, 3, 1)$	$(0, 4, 6, 8) \bmod 12$
$[3, 1, 1]$	$(1, 1, 0, 0, 1)$	$(0, 0, 1, 2) \bmod 3$
$[2, 2, 1]$	$(2, 0, 1, 1, 0)$	$(0, 0, 2, 2) \bmod 4$
$[2, 1, 1, 1]$	$(2, 1, 0, 0, 1)$	$(0, 0, 0, 2) \bmod 4$

and, second, for the Weyl group of A_5 ,

Conjugacy class $[w]$	Gradation s_w	Grades of $\mathcal{H}[w]$
$[5, 1]$	$(1, 1, 1, 0, 1, 1)$	$(0, 1, 2, 3, 4) \bmod 5$
$[4, 2]$	$(2, 1, 1, 2, 1, 1)$	$(0, 2, 4, 4, 6) \bmod 8$
$[4, 1, 1]$	$(2, 2, 1, 0, 1, 2)$	$(0, 0, 2, 4, 6) \bmod 8$
$[3, 3]$	$(1, 0, 1, 0, 1, 0)$	$(0, 1, 1, 2, 2) \bmod 3$
$[3, 2, 1]$	$(4, 1, 3, 0, 3, 1)$	$(0, 0, 4, 6, 8) \bmod 12$
$[3, 1, 1, 1]$	$(1, 1, 0, 0, 0, 1)$	$(0, 0, 0, 1, 2) \bmod 3$
$[2, 2, 2]$	$(1, 0, 0, 1, 0, 0)$	$(0, 0, 1, 1, 1) \bmod 2$
$[2, 2, 1, 1]$	$(2, 0, 1, 0, 1, 0)$	$(0, 0, 0, 2, 2) \bmod 4$
$[2, 1, 1, 1, 1,]$	$(2, 1, 0, 0, 0, 1)$	$(0, 0, 0, 0, 2) \bmod 4$

Acknowledgements

We would like to thank Tim Hollowood, Luiz Ferreira and Philippe Ruelle for their very useful comments and discussions. We are specially indebted to Laszlo Fehér for having suggested the use of lemma 9 in section 6. The research reported in this paper has been supported partially by C.I.C.Y.T (AEN93-0729) and D.G.I.C.Y.T. (PB90-0772). CRFP and MVG are also supported partly by Xunta de Galicia.

References

- [1] P. Bouwknegt and K. Schoutens, Phys. Rep. **223** (1993) 183;
P. van Driel, *Construction and Applications of Extended Conformal Symmetries*, PhD thesis, Univ. of Amsterdam (1992);
T. Tjin, *Finite and Infinite W Algebras and their Applications*, PhD thesis, Univ. of Amsterdam (1993).
- [2] A.B. Zamolodchikov, Theor. Math. Phys. **65** (1985) 347.
- [3] I.M. Gel'fand and L.A. Dikii, Funkts. Anal. Pril. **10** (1976) 13; Funkts. Anal. Pril. **13** (1979) 13.

- [4] V.G. Drinfel'd and V.V. Sokolov, J. Sov. Math. **30** (1985) 1975; Soviet. Math. Dokl. **23** (1981) 457.
- [5] S.L. Luk'yanov and V.A. Fateev, Sov. Sci. Rev. A. Phys. **15** (1990) 1.
- [6] F.A. Bais, P. Bouwknegt, K. Schoutens, and M. Surridge, Nucl. Phys. **B304** (1988) 348-370; Nucl. Phys. **B304** (1988) 371-391.
- [7] I. Bakas, Commun. Math. Phys. **123** (1989) 627; Phys. Lett. **B213** (1988) 313; P. Mathieu, Phys. Lett. **B208** (1988) 101.
- [8] F.A. Bais, T. Tjin, and P. van Driel, Nucl. Phys. **B357** (1991) 632; L. Fehér, L. O'Raifeartaigh, P. Ruelle, I. Tsutsui, and A. Wipf, Ann. Phys. (N.Y.) **213** (1992) 1.
- [9] J. Balog, L. Fehér, P. Forgács, L. O'Raifeartaigh, and A. Wipf, Ann. Phys. (N.Y.) **203** (1990) 76.
- [10] L. Fehér, L. O'Raifeartaigh, P. Ruelle, I. Tsutsui, and A. Wipf, Phys. Rep. **222** (1992) 1.
- [11] L. Frappat, E. Ragoucy and P. Sorba, Commun. Math. Phys. **157** (1993) 499.
- [12] L. Fehér, L. O'Raifeartaigh, P. Ruelle, and I. Tsutsui, Commun. Math. Phys. **162** (1994) 399.
- [13] T. Tjin and P. van Driel, *Coupled WZNW-Toda Models and Covariant KdV Hierarchies*, preprint ITFA-91-04; F. Toppan, Phys. Lett. **B327** (1994) 249.
- [14] P. Mathieu and W. Oevel, Mod. Phys. Lett. **A6** (1991) 2397.
- [15] D.A. Depireux and P. Mathieu, Int. J. of Mod. Phys. **A7** (1992) 6053.
- [16] I. Bakas and D.A. Depireux, Mod. Phys. Lett. **A6** (1991) 1561, ERRATUM *ibid.* **A6** (1991) 2351; Int. J. Mod. Phys. **A7** (1992) 1767.
- [17] P. van Driel, Phys. Lett. **274B** (1991) 179.
- [18] I. Bakas and D.A. Depireux, *Self-Duality, KdV Flows and W-Algebras*, Proceedings of the XX^{th} International Conference on Differential Geometric Methods in Theoretical Physics, New York, June 1991.
- [19] L. Fehér, J. Harnad, and I. Marshall, Commun. Math. Phys. **154** (1993) 181; L. Fehér, *Generalized Drinfeld-Sokolov Hierarchies and $\mathcal{W}$ -algebras*, Proceedings of the NSERC-CAP Workshop on Quantum Groups, Integrable Models and Statistical Systems, Kingston, Canada, 1992.
- [20] L. Bonora and C.S. Xiong, *The $(N, M)^{th}$ KdV Hierarchy and the associated W Algebras*, SISSA 171/93/EP, BONN-HE/45/93, hep-th 9311070; L. Bonora, Q.P. Liu, and C.S. Xiong, *The Integrable Hierarchy constructed from a pair of higher KdV Hierarchies and its associated W Algebra*, BONN-TH-94-17, hep-th 9408035.
- [21] M.F. de Groot, T.J. Hollowood, and J.L. Miramontes, Commun. Math. Phys. **145** (1992) 57.
- [22] N.J. Burroughs, M.F. de Groot, T.J. Hollowood, and J.L. Miramontes, Commun. Math. Phys. **153** (1993) 187; Phys. Lett. **B277** (1992) 89.

- [23] N.J. Burroughs, *Nonlinearity* **6** (1993) 583.
- [24] N.J. Burroughs, *Nucl. Phys.* **B379** (1992) 340.
- [25] F. Delduc and L. Fehér, *Conjugacy Classes in the Weyl Group Admitting a Regular Eigenvector and Integrable Hierarchies*, ENSLAPP-L-493/94, hep-th 9410203.
- [26] T.J. Hollowood, and J.L. Miramontes, *Commun. Math. Phys.* **157** (1993) 99.
- [27] V.G. Kac and M. Wakimoto, *Proceedings of Symposia in Pure Mathematics*, vol. **49** (1989) Part I, 191.
- [28] B. Kostant, *Amer. J. Math.* **81** (1959) 973.
- [29] J. Underwood, *Aspects of Non-Abelian Toda Theories*, preprint Imperial/TP/92-93/30, hep-th 9304156.
- [30] B. Feigin and E. Frenkel, *Kac-Moody Groups and Integrability of Soliton Equations*, preprint RIMS-970 (1994).
- [31] V.G. Kac and D.H. Peterson, *Symposium on Anomalies, Geometry and Topology*, W.A. Bardeen and A.R. White (eds.), Singapore, World Scientific (1985) 276-298.
- [32] N. Jacobson, *Lie Algebras*, Wiley-Interscience, New York (1962).
- [33] F. Delduc, E. Ragoucy and P. Sorba, *Phys. Lett.* **B279** (1992) 319.
- [34] V.G. Kac, *Infinite Dimensional Lie Algebras* (3rd ed.), Cambridge University Press, Cambridge (1990).
- [35] R.W. Carter and G.B. Elkington, *J. of Algebra* **20** (1972) 350.
- [36] E.B. Dynkin, *Amer. Math. Soc. Trans. Ser. 2* **6** (1957) 111.
- [37] M. Lorente and B. Gruber, *J. Math. Phys.* **13** (1972) 1639.
- [38] E. Ragoucy, *Nucl. Phys.* **B409** (1993) 213-236.
- [39] R.W. Carter, *Compositio Mathematica* **25** Fasc. 1 (1972) 1.
- [40] F. ten Kroode and J. van de Leur, *Commun. Math. Phys.* **137** (1991) 67.
- [41] F. ten Kroode and J. van de Leur, *Commun. in Algebra* **20** (1992) 3119; *Acta Appl. Math.* **27** (1992) 153; *Acta Appl. Math.* **31** (1993) 1.
- [42] C.R. Fernández-Pousa and J.L. Miramontes, in preparation.
- [43] L. Fehér, L. O’Raifeartaigh, P. Ruelle, and I. Tsutsui, *Phys. Lett.* **B283** (1992) 243.
- [44] L. Fehér, private communication.
- [45] M. Bershadsky, *Commun. Math. Phys.* **139** (1991) 71;
A. Polyakov, *Int. J. Mod. Phys.* **A5** (1990) 833.
- [46] G.W. Wilson, *Ergod. Theor. & Dyn. Sist.* **1** (1981) 361.